



**INTERNATIONAL JOURNAL OF
PHARMACEUTICAL SCIENCES**
[ISSN: 0975-4725; CODEN(USA): IJPS00]
Journal Homepage: <https://www.ijpsjournal.com>



Review Paper

Analgesic And Anti-Inflammatory Potential of Hygrophila Spinosa and Curcuma Longa: An Updated Review

Sushant Mishra*, Dhruva Kumar Yadav, Dr. Anurag Kumar

Department of Pharmacology, Shree Krishna College of Pharmacy, Sitapur, Uttar Pradesh (261125).

ARTICLE INFO

Published: 23 June 2026

Keywords:

Hygrophila Spinosa,
Curcuma longa, Analgesic,
Anti-inflammatory,
Medicinal plants,
Phytochemistry

DOI:

10.5281/zenodo.20813408

ABSTRACT

Inflammation and pain have a substantial impact on quality of life and are important causes of chronic diseases. Non-steroidal anti-inflammatory drugs (NSAIDs) and opioids continue to be the mainstay of treatment, although prolonged use of these medications is linked to severe consequences relating to the gastrointestinal tract, kidneys, heart, and dependency. As a result, medicinal plants with anti-inflammatory and analgesic qualities have drawn more and more interest from scientists. Hygrophila Spinosa (syn. Curcuma longa and Hygrophila auriculata) have long been used in Ayurvedic medicine to treat rheumatism, pain, inflammation, and other conditions. Flavonoids, alkaloids, terpenoids, sterols, phenolic chemicals, and curcuminoids, which are responsible for a variety of pharmacological effects, have been identified via phytochemical studies. By modifying cyclooxygenase pathways, inflammatory cytokines, oxidative stress, and nociceptive mediators, experimental investigations show notable analgesic and anti-inflammatory benefits. This review summarizes the botanical aspects, phytochemistry, pharmacological evidence, mechanisms of action, safety profiles, and future therapeutic prospects of these medicinal plants, highlighting their potential as alternative or adjunctive therapies for inflammatory disorders.

INTRODUCTION

The body uses inflammation as a defence mechanism against viruses, tissue damage, and toxic substances. On the other hand, chronic inflammation has a role in the development of cancer, diabetes, neurological illnesses, cardiovascular disease, and arthritis. One of the most frequent reasons for seeking medical

attention is pain, which is intimately linked to inflammatory processes.

NSAIDs and corticosteroids are examples of conventional anti-inflammatory drugs that are helpful but can have negative side effects such as gastrointestinal ulcers, nephrotoxicity, and cardiovascular problems. As a result, medicinal plants with enhanced safety profiles and anti-

*Corresponding Author: Sushant Mishra

Address: Department of Pharmacology, Shree Krishna College of Pharmacy, Sitapur, Uttar Pradesh (261125)..

Email ✉: Shushantmishra123@gmail.com

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.



inflammatory and analgesic qualities are gaining popularity.

Hygrophila spinosa and *Curcuma longa* are two traditional medicinal plants that have garnered a lot of interest because of their widespread ethnomedical use and pharmacological properties that have been verified by science. Their phytoconstituents include immunomodulatory, anti-inflammatory, antioxidant, and antinociceptive qualities, indicating possible therapeutic uses in inflammatory conditions.

2. METHODOLOGY OF LITERATURE SEARCH

The literature was collected from:

- PubMed
- Scopus
- Web of Science
- Google Scholar
- ScienceDirect

Publications from 2000–2025 were reviewed with emphasis on peer-reviewed research articles, review papers, and pharmacological studies.

3. BOTANICAL PROFILE OF HYGROPHILA SPINOSA TAXONOMY

Parameter	= Description
Family =	Acanthaceae
Genus =	Hygrophila
Species =	Hygrophila spinosa T. Anders
Synonym =	Hygrophila auriculata

TRADITIONAL USES

- Rheumatism
- Joint pain
- Inflammation
- Urinary disorders
- Hepatic disorders
- Edema
- General debility

The plant contains flavonoids, alkaloids, sterols, terpenoids, tannins, and phenolic compounds contributing to its medicinal properties

MAJOR PHYTOCHEMICAL CLASSES

- Alkaloids
- Flavonoids
- Steroids
- Triterpenoids
- Saponins
- Tannins
- Glycosides
- Phenolic compounds



FIGURE 1. BOTANICAL REPRESENTATION OF *HYGROPHILA SPINOSA*

TABLE 1. MAJOR PHYTOCHEMICAL CONSTITUENTS OF *HYGROPHILA SPINOSA*

Phytochemicals Tested	Plant extracts of <i>Hygrophila auriculata</i>				
	Aqueous	Acetone	Chloroform	Petroleum ether	Ethanol
Tannins	++	++	+	-	+
Saponins	++	+	+	+	-
Quinones	+	++	+	-	++
Terpenoids	+	++	+	+	++
Steroids	+	+	+	+	++
Flavonoids	++	++	+	-	+
Phenol	++	++	++	+	++
Alkaloids	-	-	-	-	+
Glycosides	-	-	-	-	-
Cardiac glycosides	-	++	+	-	+
Coumarins	+	+	+	-	+
Antho cyanin	-	-	-	-	-
Beta cyanin	++	+	+	-	+

++ = strong positive; + = positive; - = negative

PATHOPHYSIOLOGY OF PAIN AND INFLAMMATION AND THE ROLE OF *HYGROPHILA SPINOSA*

After tissue damage, infection, or exposure to damaging stimuli, pain and inflammation are closely associated physiological reactions. Redness, swelling, heat, pain, and loss of function are signs of inflammation. Inflammatory mediators like histamine, serotonin, bradykinin, prostaglandins, leukotrienes, and pro-inflammatory cytokines like tumor necrosis factor- α (TNF- α), interleukin-1 β (IL-1 β), and interleukin-6 (IL-6) are released when immune cells like neutrophils, mast cells, and macrophages are activated.

These mediators encourage leukocyte migration to the site of damage by increasing vascular permeability. Concurrently, prostaglandins produced by the cyclooxygenase (COX) pathway cause nociceptors to become more sensitive, which causes pain perception. By continuously activating peripheral and central sensitization processes, persistent inflammation can result in chronic pain.

Numerous bioactive phytochemicals, such as flavonoids, alkaloids, sterols, triterpenoids, tannins, and phenolic compounds, are present in *Hygrophila Spinosa*. Through a variety of

processes, these components support its analgesic and anti-inflammatory properties. By scavenging reactive oxygen species (ROS) and lowering oxidative stress, which is a significant factor in inflammatory tissue damage, flavonoids and phenolic compounds function as powerful antioxidants.

Experimental studies suggest that extracts of *H. spinosa* inhibit the synthesis and release of inflammatory mediators such as prostaglandins and cytokines. The plant may suppress the activation of nuclear factor-kappa B (NF- κ B), a key transcription factor involved in the expression of inflammatory genes. Inhibition of NF- κ B leads to reduced production of TNF- α , IL-1 β , and IL-6, thereby attenuating the inflammatory response.

The analgesic effect of *H. spinosa* is attributed to its ability to interfere with peripheral pain pathways by reducing prostaglandin formation and inflammatory mediator release. Additionally, certain phytoconstituents may exert central analgesic effects by modulating neurotransmitter systems involved in pain transmission. Reduction of oxidative stress and inflammatory cytokine production further contributes to pain relief.

Overall, the anti-inflammatory and analgesic properties of *Hygrophila spinosa* are mediated through antioxidant activity, inhibition of



inflammatory mediator synthesis, suppression of pro-inflammatory cytokines, stabilization of cellular membranes, and modulation of pain signaling pathways. These pharmacological actions support its traditional use in the management of inflammatory disorders and painful conditions.

4. BOTANICAL PROFILE OF *CURCUMA LONGA*

TAXONOMY

PARAMETER	DESCRIPTION
Family =	Zingiberaceae
Genus =	Curcuma
Species=	Curcuma longa Linn.

TRADITIONAL USES

- Arthritis
- Wound healing
- Pain management
- Inflammatory diseases
- Gastrointestinal disorders

The rhizome is rich in curcuminoids and volatile oils responsible for therapeutic activity. Another important factor in the pathophysiology of inflammation is oxidative stress. Reactive oxygen species (ROS) and reactive nitrogen species (RNS) produced by activated immune cells during inflammation result in lipid peroxidation, cellular damage, and a



FIGURE 2. BOTANICAL REPRESENTATION OF *CURCUMA LONGA*:

5. PHYTOCHEMISTRY OF *HYGROPHILA SPINOSA*

Major phytoconstituents include:

FLAVONOIDS

- Apigenin
- Luteolin
- Quercetin derivatives

TRITERPENOIDS

- Lupeol
- Betulin
- Lupenone

STEROLS

- β -Sitosterol
- Stigmasterol

ALKALOIDS

- Asteracanthine
- Asteracanthicine

OTHER CONSTITUENTS

- Tannins
- Phenolics
- Fatty acids
- Glycosides

These compounds exhibit antioxidant, anti-inflammatory, and analgesic effects.

6. PHYTOCHEMISTRY OF *CURCUMA LONGA*

CURCUMINOIDS

- Curcumin
- Demethoxycurcumin
- Bisdemethoxycurcumin

ESSENTIAL OILS

- Turmerone
- Atlantone
- Zingiberene

OTHER CONSTITUENTS

- Proteins
- Polysaccharides
- Phenolic compounds

Curcumin is considered the principal bioactive compound responsible for anti-inflammatory and analgesic actions.

PATHOPHYSIOLOGY OF PAIN AND INFLAMMATION AND THE ROLE OF *CURCUMA LONGA*

Inflammation and pain are defensive physiological reactions brought on by immune-mediated harm, infection, or tissue damage. When immune cells like neutrophils, macrophages, and mast cells are activated during acute inflammation, they release inflammatory mediators such histamine, serotonin, bradykinin, prostaglandins, leukotrienes, and cytokines. These mediators cause pain, redness, swelling, and heat by increasing vascular permeability, encouraging leukocyte infiltration, and stimulating nociceptors. The transcription factor Nuclear Factor Kappa B, which controls the production of pro-inflammatory genes encoding cytokines like TNF- α , IL-1 β , IL-6, cyclooxygenase-2 (COX-2), and inducible nitric oxide synthase (iNOS), is activated during inflammation. Persistent pain and chronic inflammatory disorders are linked to the overproduction of these mediators.

n increase in inflammatory signaling pathways. Pain receptor sensitization and tissue damage are encouraged by prolonged oxidative stress.

Curcumin is the main pharmacologically active polyphenol found in *Curcuma longa*, or turmeric. Through a variety of molecular pathways, curcumin demonstrates strong analgesic and anti-inflammatory properties. It decreases the transcription of inflammatory cytokines, chemokines, COX-2, and iNOS via suppressing NF- κ B activation. As a result, prostaglandins, nitric oxide, TNF- α , IL-1 β , and IL-6 are produced at much lower levels.

Additionally, curcumin reduces the production of prostaglandins and leukotrienes, which influence pain and inflammation, by inhibiting the activity

of the enzymes cyclooxygenase and lipoxygenase. Curcumin also has potent antioxidant qualities, scavenging free radicals and boosting natural antioxidant defenses including glutathione peroxidase, catalase, and superoxide dismutase (SOD). *Curcuma longa*'s analgesic effects are mainly ascribed to its regulation of pain-signaling pathways and suppression of inflammatory mediator production. By lowering prostaglandin synthesis, curcumin lessens peripheral sensitivity. It may also affect central pain pathways by modifying neurotransmitters and inflammatory signals in the central nervous system.

Recent studies further indicate that curcumin regulates multiple intracellular signaling cascades, including the mitogen-activated protein kinase (MAPK) pathway, Janus kinase/signal transducer and activator of transcription (JAK/STAT) pathway, and Toll-like receptor (TLR)-mediated inflammatory responses. Through these mechanisms, *Curcuma longa* effectively suppresses both acute and chronic inflammatory processes.

7. ANALGESIC ACTIVITY OF *HYGROPHILA SPINOSA*

Experimental studies have demonstrated significant analgesic activity using:

- Hot plate test
- Tail flick test
- Acetic acid-induced writhing test

Extracts of alcohol and chloroform dramatically decreased pain reactions, indicating both peripheral and central analgesic processes. Flavonoids, sterols, and triterpenoids that affect prostaglandin synthesis and nociceptive pathways may be responsible for the reported action.

8. ANTI-INFLAMMATORY ACTIVITY OF *HYGROPHILA SPINOSA*

Studies using carrageenan-induced paw edema models demonstrated significant inhibition of



inflammation by alcoholic and chloroform extracts.

Proposed mechanisms include:

- Inhibition of prostaglandin synthesis
- Reduction of inflammatory mediators

These findings support the traditional use of the plant in rheumatism and inflammatory disorders.

9. ANALGESIC ACTIVITY OF *CURCUMA LONGA*

Curcuma longa exhibits potent antinociceptive activity in:

- Formalin test
- Acetic acid writhing test
- Tail immersion test

Curcumin reduces pain perception by modulating:

- Substance P
- TRPV1 receptors
- Cyclooxygenase pathways
- Nitric oxide production

Experimental studies suggest efficacy comparable to standard analgesics in some models.

10. ANTI-INFLAMMATORY ACTIVITY OF *CURCUMA LONGA*

Curcumin exerts anti-inflammatory effects through multiple pathways:

SUPPRESSION OF PRO-INFLAMMATORY CYTOKINES

- TNF- α
- IL-1 β
- IL-6

INHIBITION OF ENZYMES

- COX-2
- LOX
- iNOS

MODULATION OF SIGNALING PATHWAYS

- NF- κ B
- MAPK

- JAK/STAT

These mechanisms contribute to the reduction of edema, inflammatory cell infiltration, and oxidative damage.

11. COMPARATIVE PHARMACOLOGICAL PROFILE

PARAMETER	HYGROPHILA SPINOSA	CURCUMA LONGA
Major Actives	Flavonoids, Lupeol, Sterols	Curcuminoids
Analgesic Activity	Significant	Strong
Anti-inflammatory Activity	Significant	Strong
Antioxidant Activity	High	Very High
Clinical Evidence	Limited	Extensive
Safety Profile	Good	Excellent

13. TOXICOLOGICAL ASSESSMENT

H. spinosa extracts have comparatively low toxicity at therapeutic dosages, according to acute toxicity tests. The experimental animals did not exhibit any notable behavioral problems or death. But there are still a few restrictions:

- Studies on chronic toxicity are lacking.
- Limited assessment of reproductive toxicity
- Inadequate information on human safety
- The lack of uniform dosage recommendations

Prior to clinical application, additional toxicological research is necessary.

14. FUTURE PERSPECTIVES

Future research should focus on:

1. Isolation of active constituents.
2. Standardization of extracts.
3. Pharmacokinetic studies.
4. Toxicological evaluation.
5. Clinical trials.
6. Nanoformulation development.



7. Evaluation of synergistic combinations.

CONCLUSION

According to available data, the various phytochemical components of *Hygrophila Spinosa* and *Curcuma longa* may have important analgesic and anti-inflammatory effects. Their traditional application in pain and inflammatory conditions is supported by experimental research. In contrast to *Hygrophila spinosa*, which has promising pharmacological potential but is still understudied, *Curcuma longa* has considerably stronger clinical data. For the treatment of chronic inflammatory disorders and pain syndromes, combined formulations may offer a novel therapeutic approach. To determine efficacy, safety, and standardization for therapeutic usage, more clinical research is needed.

REFERENCES

1. Indian Materia Medica. Nadkarni AK. *Indian Materia Medica*. Mumbai: Popular Prakashan; 2007.
2. Kshirsagar AD, Ingale KG, Vyawahare NS, Thorve VS. *Hygrophila spinosa*: A comprehensive review. *Pharmacognosy Reviews*. 2010;4(8):167–171.
3. Gaikwad S, Kothapalli L, Thomas A. Phytochemical and Pharmacological Aspects of *Hygrophila Spinosa*: An Overview. *Current Traditional Medicine*. 2023.
4. Patra A, Jha S, Murthy PN. Phytochemical and Pharmacological Potential of *Hygrophila spinosa* T. Anders. *Pharmacognosy Reviews*. 2009.
5. Kumar V, Abbas AK, et al. Ethnomedicinal, phytochemical and pharmacological updates on *Hygrophila auriculata*. *Journal of Integrative Medicine*. 2018.
6. Ahmed S, Rahman M, Islam M. Anti-inflammatory activity of *Hygrophila auriculata* extracts in experimental animal models.
7. Singh A, Malhotra S. Analgesic activity of *Hygrophila spinosa* leaf extracts using hot plate and writhing tests.
8. Rao KN, et al. Evaluation of anti-inflammatory potential of *Asteracantha longifolia* in carrageenan-induced paw edema.
9. Shanmugasundaram P, Venkataraman S. Hepatoprotective and antioxidant activities of *Hygrophila auriculata*.
10. Fuloria S, Mehta J, Chandel A, et al. A Comprehensive Review on the Therapeutic Potential of *Curcuma longa* Linn. in Relation to its Major Active Constituent Curcumin. *Frontiers in Pharmacology*. 2022.
11. Rahaman MM, Rakib A, Mitra S, et al. The Genus *Curcuma* and Inflammation: Overview of the Pharmacological Perspectives. *Plants*. 2021;10(1):63.
12. Amalraj A, Pius A, Gopi S, Gopi S. Biological activities of curcuminoids and turmeric. *Journal of Traditional and Complementary Medicine*. 2017.
13. Hewlings SJ, Kalman DS. Curcumin: A Review of Its Effects on Human Health. *Foods*. 2017.
14. Gupta SC, Patchva S, Aggarwal BB. Therapeutic Roles of Curcumin: Lessons Learned from Clinical Trials. *AAPS Journal*. 2013.
15. Aggarwal BB, Harikumar KB. Potential Therapeutic Effects of Curcumin. *International Journal of Biochemistry & Cell Biology*. 2009.
16. Menon VP, Sudheer AR. Antioxidant and Anti-inflammatory Properties of Curcumin. *Advances in Experimental Medicine and Biology*. 2007.
17. Joe B, Vijaykumar M, Lokesh BR. Biological Properties of Curcumin. *Critical Reviews in Food Science and Nutrition*. 2004.



18. Chainani-Wu N. Safety and Anti-inflammatory Activity of Curcumin. *Alternative Medicine Review*. 2003.
19. Jurenka JS. Anti-inflammatory Properties of Curcumin. *Alternative Medicine Review*. 2009.
20. Goel A, Kunnumakkara AB, Aggarwal BB. Curcumin as “Curecumin”: From Kitchen to Clinic. *Biochemical Pharmacology*. 2008.

HOW TO CITE: Sushant Mishra, Dhruva Kumar Yadav, Dr. Anurag Kumar, Analgesic And Anti-Inflammatory Potential of *Hygrophila Spinosa* and *Curcuma Longa*: An Updated Review, *Int. J. of Pharm. Sci.*, 2026, Vol 4, Issue 6, 5959-5966, <https://doi.org/10.5281/zenodo.20813408>

