



**INTERNATIONAL JOURNAL OF
PHARMACEUTICAL SCIENCES**
[ISSN: 0975-4725; CODEN(USA): IJPS00]
Journal Homepage: <https://www.ijpsjournal.com>



Research Paper

Antioxidant And Organprotective Activities of Amaranthus Gangeticus Leaves Against Paracetamol Induced Hepatic Damage in Albino Rats

Jyoti Jain^{*1}, Chandrika Kampli²

¹Dept. of Pharmacology, Prof. & H.O.D. MAM College of Pharmacy, Gulbarga, (Karnataka) India.

²Medical co-ordinator, Al badar Dental College & Hospital Gulbarga (Karnataka).

ARTICLE INFO

Published: 24 Aug 2026

Keywords:

Amaranthus gangeticus leaves, paracetamol, serum markers and polyphenols.

DOI:

10.5281/zenodo.22076150

ABSTRACT

Amaranthus gangeticus is a medicinal plant. Our aim is to investigate its organ protection property. Ethanolic extract of Amaranthus gangeticus leaves was taken and the parameters studied were serum glutamate pyruvate transaminase, serum glutamate oxaloacetate transaminase, total bilirubin, direct bilirubin, total cholesterol, high density lipoproteins and alkaline phosphatase activities. Bio-chemical studies of blood samples of paracetamol treated animals showed significant increase in the levels of serum markers and decrease in high density lipoproteins reflecting the liver injury caused by paracetamol. Whereas, the animals treated with ethanolic extract showed significant dose dependent decrease in the elevated levels of serum markers and increase in high density lipoproteins indicating the protection of hepatic cells. Results were analyzed by one-way analysis of variance by Dunnet's 't' test. Therefore ethanolic extract afford significant protection against paracetamol induced hepatocellular injury and remarkable rejuvenation of these tissues found in histopathological studies which may be due to polyphenols and its antioxidant activities.

INTRODUCTION

Liver is an important organ in our body. It is one of the vital organ of metabolism and excretion. During its normal physiological functioning it metabolizes various endogenous and exogenously administered chemicals, so as to terminate or inactivate these agents. Hence due to this function,

it protects the whole body from the various environmental and chemical challenges. In addition to this liver has got an inbuilt mechanism to protect itself and to regenerate on several occasions, many of these hepatotoxic challenges overpower inbuilt protective mechanism and cause hepatotoxicity resulting in the hepatic necrosis and hepatitis. *Amaranthus gangeticus* is a edible plant

***Corresponding Author:** Jyoti Jain

Address: Dept. of Pharmacology, Prof. & H.O.D. MAM College of Pharmacy, Gulbarga, (Karnataka) India.

Email ✉: jyotilcp2@gmail.com

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.



used as vegetable, it is also used by native practitioner as hepatoprotective in treating various types of jaundice. The leaves of this plant contain polyphenolic compounds like tannins and flavonoids. These polyphenolic compounds have antioxidant property and anti-oxidants have known to possess hepatoprotective activity. Keeping the native knowledge and the above mentioned literature information¹, this plant was selected for present study to screen the leaves of this edible plant for the presence of phytoconstituents, antioxidant and hepatoprotective activities. This study was carried out by using 90% ethanolic extract of *Amaranthus gangeticus* (AG) as hepatoprotectant and paracetamol (PCM) as hepatotoxicant.

MATERIALS AND METHODS

Collection and identification of plant: The plant was collected from Kusnoor village (Gulbarga district), Karnataka in the month of March and was authenticated by Dr. Srinath Rao, chairman, P.G. Department of Studies and Research in Botany, Gulbarga University, Gulbarga, Karnataka. The plant was thoroughly cleaned to remove adherent soil and other impurities, the leaves were shade dried and made into a coarse powder by rubbing in the palms.

Extraction

90 gms of shade dried leaf powder of AG was extracted in Soxhlet's apparatus using petroleum ether for defatting and then it was extracted with 90% ethanol (AGEE). The solvent was evaporated on a water bath at a low temperature (50°C) and finally the residue was obtained.

Materials used

Paracetamol (Esteem Pharmaceutical Pvt. Ltd. Agra), Silymarin (SD fine chemicals, Mumbai), Ready to use diagnostic kits (Aspen Labs Pvt. Ltd.,

Delhi-India), 90% ethanolic extract of AG, gallic acid (Rolex Chemical Industries, Mumbai). All chemicals and reagents used were of analytical grade.

Animals used

Wistar albino rats of either sex weighing between 150-180 gms were housed in polypropylene cages and were maintained at 27° ± 2°C with 12:12 hr, light/dark cycle. They were fed with commercial diet (VRK Nutritional Laboratory, Sangli) and water *ad libitum*, during the experiment. The study was permitted by Institutional Animal Ethical Committee (Reg. No. 342).

I. Evaluation of hepatoprotective activity in PCM-induced hepatotoxicity:

The method reported by R.R. Chattopadhyay was followed². In the dose response experiment, albino wistar rats were randomly assigned into 5 groups of 6 animals in each.

Gp-I Animals (-ve control) were administered with 1ml/kg p.o. of saline for 7 days. Gp-II Animals (+ve control) were administered with 1ml/kg p.o. of saline for 7 days. Gp-III Animals were administered with silymarin 100mg/kg p.o., for 7 days. Gp-IV Animals were administered with AGEe 200mg/kg p.o., for 7 days.

Gp-V Animals were administered with AGEe 400mg/kg p.o., for 7 days. On 5th day, 30 minutes after the administration of normal saline, 100mg/kg silymarin, 200mg/kg and 400mg/kg of AGEe to Group-II, III, IV and V respectively, Paracetamol 2gm/kg was given orally. After 48 hours of paracetamol dosing, rats were sacrificed under mild ether anaesthesia and the blood samples were collected from the animals of each group through carotoid artery puncture and centrifuged immediately to get clear serum, for evaluating the serum biochemical parameters by using Aspen diagnostic kits. Liver was dissected



out, the blood was blotted off, washed with saline and stored in 10% formalin and proceeded for histopathology to evaluate the details of hepatic architecture in each group microscopically. The results are shown in table-1.

II. Total Phenolic Content estimation³

1. Extraction of phenols from *Amaranthus gangeticus* leaves⁴

Method: In this process, 10 gms of shade dried, leaf powder of *Amaranthus gangeticus* (AG) is placed with the whole of the menstrum (250 ml) in a closed vessel for 2 days. During this period shaking is done occasionally. After 2 days, the liquid is strained and marc is pressed. The expressed liquid is mixed with strained liquid. It is then filtered to make a clear liquid. The final volume is not adjusted. It is evaporated on a water-bath at low temperature not exceeding (50°C) and preserved. Plant extracts were prepared using two different extracting solvents :

AG_{CW} : *Amaranthus gangeticus* shade dried leaf powder macerated for 48 hrs with cold water (distilled water stored at room temperature).

AG_{HW} : AG shade dried leaf powder macerated for 48 hrs with hot water (50°C).

Total Phenolic Content

It was estimated by using Folin-Ciocalteu reagent according to the method reported by Singleton and Rossi³ using standard curve generated with Gallic acid.

Procedure: For the preparation of Calibration curve, a series of calibrated 10 ml volumetric flasks were taken and appropriate aliquots of the working standard solution of gallic acid were pipetted out. To each flask, 5ml of Folin-Ciocalteu reagent (diluted ten fold) and 4ml of sodium carbonate solution (75g/l) was mixed with appropriate aliquots of gallic acid. The absorbance was measured after 30 min. at 765nm in Shimadzu 1700 UV-visible spectrophotometer. Same procedure was applied for various extracts of *Amaranthus gangeticus* (1mg/ml) was mixed with the same reagent as described in the construction of calibration curve and after 30 min. the absorbance was measured for the determination of total phenolic compound. The total phenolic content was expressed as gallic acid equivalents (GAE) in milligram per gram of sample using a standard curve of gallic acid. Total amount of phenolic compounds in *Amaranthus gangeticus* were done in triplicates and concentrations of total phenolic content in various extracts were determined and expressed as gallic acid in mg equivalents per gm of sample using the standard curve generated with gallic acid. The results obtained were compiled and represented as shown in table-2.

Statistical analysis

The data presented in table-1 (n=6) and 2 (n=3) were expressed as mean $\pm$ standard error of mean (SEM). Significant difference among the mean were calculated at the level of $p < 0.001$ and analyzed by one-way analysis of variance by Dunnet's 't' test. A value of $p < 0.05$ was defined as significant.



Table-1 Effect of 90% ethanolic extract of *Amaranthus Gangeticus* leaves on hepatic enzymes in PCM induced hepatotoxicity

Treatment	Biochemical parameters Mean $\pm$ SEM						
	SGPT IU/L	SGOT IU/L	Total Bilirubin mg/dl	Direct Bilirubin mg/dl	Total Cholesterol mg/dl	HDL mg/dl	ALP IU/L
Negative control (1ml/ kg saline p.o.)	63.59 $\pm$ 0.33	71.91 $\pm$ 0.69	0.95 $\pm$ 0.008	0.27 $\pm$ 0.001	117.6 $\pm$ 0.76	8.18 $\pm$ 0.005	138 $\pm$ 0.8
PCM treated (positive control) (1ml/kg saline p.o. + 2gms/kg p.o.)	206.5 $\pm$ 0.57	270.55 $\pm$ 0.58	4.12 $\pm$ 0.006	0.90 $\pm$ 0.005	195.6 $\pm$ 0.71	4.35 $\pm$ 0.007	339.5 $\pm$ 0.75
Silymarin+ PCM (100mg/kg p.o.+ 2gms /kg p.o.)	75.92 $\pm$ 0.45***	105.1 $\pm$ 0.78***	1.16 $\pm$ 0.006***	0.32 $\pm$ 0.001***	115.8 $\pm$ 0.47***	7.09 $\pm$ 0.007***	148.5 $\pm$ 0.89***
AGEE + PCM (200mg/kg p.o. + 2gms /kg, p.o)	162.3 $\pm$ 0.76***	175.3 $\pm$ 0.57***	2.34 $\pm$ 0.005***	0.52 $\pm$ 0.001***	151.5 $\pm$ 0.42***	5.06 $\pm$ 0.004***	240.0 $\pm$ 1.06***
AGEE + PCM (400mg/kg p.o. + 2gms /kg, p.o)	89 $\pm$ 0.36***	112.0 $\pm$ 0.57***	1.41 $\pm$ 0.006***	0.34 $\pm$ 0.001***	129.5 $\pm$ 0.42***	6.71 $\pm$ 0.007***	152.8 $\pm$ 0.60***

Values are the mean $\pm$ S.E.M. of six rats/treatment.

*** P<0.001 Significance compared to PCM treatment.

Table-2 Total phenol content in *amaranthus gangeticus* leaf extract

Sl. No.	Name of the extract	Total phenol content mg/gms (in gallic acid equivalent)
1.	AGEE	251.484 $\pm$ 1.92
2.	AG _{CW}	127.097 $\pm$ 1.73
3.	AG _{HW}	192.070 $\pm$ 2.63

RESULTS

There is a marked increase in serum biomarkers which were observed in PCM treated group. But, these levels were reversed to near normal levels with the treatment of 200mg/kg and 400mg/kg of 90% ethanolic extract of AG. Whereas, the standard silymarin has restored the serum biomarker levels significantly which is depicted in table-1. Results showed that, among the three different extracts of *Amaranthus gangeticus* AG_{CW} contained the lowest level of total phenol content, while AGEe contained the highest level. The observations are recorded in table-2.

DISCUSSIONS

The model selected to assess organ protection was the 90% ethanolic extract of AG which was tested against PCM induced hepatotoxicity in rats. The PCM treated group exhibited extensive fatty changes, congestion of sinusoids, necrosis etc. upon histopathological observations. But treatment with AGEe, the serum levels of biochemical markers of hepatocellular damage like SGPT, SGOT, bilirubin (total and direct), ALP, cholesterol were decreased as a mark of fatty change, congestion, inflammation etc. The animals treated with 90% ethanolic extract of AG for 7



days showed remarkable rejuvenation of hepatocellular architecture, the AGEe reversed the elevated serum markers of liver damage proportionate to the doses employed. Protection was offered by silymarin, (100mg/kg), 200mg/kg and 400mg/kg of AGEe respectively (compared to positive control w.r.t. biochemical parameters). The hepatoprotection offered by AG may be attributed due to the presence of antioxidant phyto-constituents like flavonoids phytosterols and other polyphenolic constituents, by the virtue of which the extract nullified the powerful hepatotoxic radicals generated by PCM (i.e., hydroxyl radical and reactive oxygen species)⁵⁻⁷ before they could initiate fatal consequences. These findings add strength to our claim. Total Phenolic Content of AGEe, AG_{CW} and AG_{HW} were determined. Total phenolic content was maximum in AGEe (251.484 mg/gms in terms of gallic acid equivalent).

CONCLUSIONS

90% Ethanolic extract of AG has a powerful organ protection property and it also has a good *in-vitro* antioxidant property which are attributed due to presence of antioxidant phyto-constituents. Therefore, the above findings reveals that the use of *Amaranthus gangeticus* leaves in our food protects our liver.

SCOPE FOR FUTURE STUDY

As it is a medicinal plant, hence isolation of its phytoconstituents are needed to screen various organ protective potentials.

ACKNOWLEDGEMENTS

We are thankful to the Management and Principal of MAM college of Pharmacy, Gulbarga for providing all the necessary facilities to carry out this research work.

CONFLICT OF INTEREST

The author declares no conflict of interest.

REFERENCES

1. www.google.co.in – wikipedia, the free encyclopedia : navigation search
2. Chattopadhyay RR, Possible mechanism of hepatoprotective activity of *Azadirachta indica* leaf extract: Part II J. Ethnopharmacol 2003; 89: 217-19.
3. Singleton V.L. and Rossi J.A. (1965). Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. Am. J. Enol vit. c, 16, 144-58.
4. Mehta RM, Pharmaceutics-I, 4th ed., Vallabh Prakashan, Delhi, 2007. pp. 149.
5. Attri S, Rana SV, Vaiphei K, Sodhi CP, Katyal R, Nain RC, Goel K, Singh K. Isoniazid and rifampicin-induced oxidative hepatic injury-protection by N-acetylcysteine. Hum. Exp. Toxicol. 2000; 19: 517-22.
6. Sadanobu S, Watanabe M, Nakamura C, Tezuka M. In vitro tests of 1,3-dithia-2thioxo-cyclopent-4-ene to evaluate the mechanisms of its hepatoprotective action. J. Toxicol. Sci. 1999; 24: 375-81.
7. Ramirez D, Commandeur JNM, Groot E, Vermeulen NPE. Mechanism of protection of lobenzarit against paracetamol-induced toxicity in rat hepatocytes. Eur. J. Pharmacol. 1995; 293:301-

HOW TO CITE: Jyoti Jain, Chandrika Kampli, Antioxidant And Organprotective Activities of *Amaranthus Gangeticus* Leaves Against Paracetamol Induced Hepatic Damage in Albino Rats, Int. J. of Pharm. Sci., 2026, Vol 4, Issue 8, 3779-3783, <https://doi.org/10.5281/zenodo.22076150>

