



**INTERNATIONAL JOURNAL OF
PHARMACEUTICAL SCIENCES**
[ISSN: 0975-4725; CODEN(USA): IJPS00]
Journal Homepage: <https://www.ijpsjournal.com>



Research Article

Evaluation of Anti Urolethiatic Potential of Vitis Vinifera Leaves

Sumit Shinge¹, Sumayya Shaikh², Sujit Chavan³, Sujit Ukande⁴, Sushant Patil⁵

Dr. J.J. Magdum Pharmacy Collage, Jaysingpur

ARTICLE INFO

Published: 25 Jun. 2026

Keywords:

Urolithiasis; Kidney stones;
Medicinal plants;
Antirolithiatic activity;
Calcium oxalate;
Phytochemicals; Oxidative
stress; Nephroprotection;
Herbal therapy

DOI:

10.5281/zenodo.20844672

ABSTRACT

Urolithiasis is a common urinary tract disorder characterized by the formation of kidney stones due to the crystallization of minerals in the urinary system. The present study was carried out to evaluate the anti-urolithiatic activity of Vitis vinifera leaves extract using an in-vitro nucleation assay method. The leaves of Vitis vinifera were collected, dried, powdered, and extracted using ethanol by Soxhlet extraction method. Preliminary phytochemical screening of the extract revealed the presence of flavonoids, tannins, phenolic compounds, saponins, and alkaloids, which are known for their medicinal properties. The anti-urolithiatic activity was assessed by measuring the inhibition of calcium oxalate crystal nucleation at different concentrations of the extract and comparing it with the standard drug Cystone. Absorbance values were recorded using a UV-Visible spectrophotometer, and percentage inhibition was calculated. The extract showed concentration-dependent inhibition of calcium oxalate crystal formation, indicating significant anti-urolithiatic potential. Higher concentrations of the extract exhibited greater inhibitory activity, comparable to the standard drug. The study suggests that Vitis vinifera leaves possess promising anti-urolithiatic activity and may be useful in the prevention and management of kidney stone formation. Further in-vivo and clinical studies are required to confirm its therapeutic efficacy and safety.

INTRODUCTION

The kidneys are two bean-shaped organs, each approximately the size of a fist, located on either side of the spine just below the rib cage. They play a vital role in maintaining overall body health by filtering waste products, toxins, and excess fluids from the blood to produce urine. In addition to waste removal, the kidneys help regulate

electrolyte balance, maintain fluid levels, control blood pressure, and support the production of red blood cells through hormone secretion. Their proper functioning is essential for maintaining the body's internal balance and overall well-being. [1]

Urolithiasis is a common disorder of the urinary system characterized by the formation of stones (calculi) within the kidneys, ureters, bladder, or

***Corresponding Author:** Sumit Shinge

Address: Dr. J.J. Magdum Pharmacy Collage, Jaysingpur.

Email ✉: sumitshinge405@gmail.com

Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.



urethra. The term is derived from the Greek words ouron (urine) and lithos (stone), referring to the formation of urinary stones. Among the various types of urinary calculi, calcium oxalate stones are the most prevalent. These stones develop when calcium ions combine with oxalate in the urine, leading to the formation and aggregation of crystals. Over time, these crystals grow into larger stone masses, which can cause pain, urinary obstruction, and other complications if left untreated.[2]

Nephrolithiasis, commonly known as kidney stone disease, is a significant public health concern worldwide. In India, approximately two million people are affected by kidney stones each year. Certain regions, including Gujarat, Maharashtra, Punjab, Rajasthan, Delhi, Haryana, and parts of northern India, are commonly referred to as the "stone belt" due to the high prevalence of the disease. Kidney stones are also frequently reported in southern India, where dietary habits, including the regular consumption of oxalate-rich foods such as tamarind, may contribute to stone formation.

Globally, the incidence and prevalence of kidney stones vary across different regions. Studies indicate that approximately 0.1–0.4% of the population in the United States and Europe develops kidney stones annually. The lifetime prevalence is estimated to be 2–5% in Asia, 8–15% in Europe and North America, and up to 20% in Saudi Arabia. Kidney stone disease is also known for its high recurrence rate, with nearly 75% of affected individuals experiencing a recurrence within 20 years of the initial episode. These statistics highlight the growing burden of urolithiasis and the importance of effective preventive and therapeutic strategies.[3]

The treatment of kidney stones depends on the size, type, and location of the stone as well as the severity of symptoms. Small stones are usually

treated conservatively by increasing fluid intake to 2–3 liters per day, which helps dilute urine and promotes stone passage. Large or obstructive stones require surgical or minimally invasive procedures such as Extracorporeal Shock Wave Lithotripsy, which uses shock waves to break stones into smaller fragments, Ureteroscopy, where a scope is inserted through the urinary tract to remove or fragment the stone, and Percutaneous Nephrolithotomy, which is used for large kidney stones through a small incision in the back.[4]

Synthetic drug used in treatment of Kidney Stones involves the use of chemically synthesized drugs and modern medical procedures to relieve pain, dissolve stones, prevent crystal formation, and promote stone expulsion. The first step in treatment is pain management because kidney stones cause severe flank pain due to obstruction of urine flow. Non-steroidal anti-inflammatory drugs such as Ibuprofen and Diclofenac are commonly used to reduce pain and inflammation, while stronger analgesics may be given in severe cases.

Drugs like Tamsulosin and Nifedipine relax the smooth muscles of the ureter and facilitate the passage of stones through urine. In patients with uric acid stones, alkalinizing agents such as potassium citrate and sodium bicarbonate are administered to increase urinary pH and dissolve the stones. Allopurinol is prescribed to reduce uric acid production and prevent recurrence of uric acid stones. For calcium stone prevention, thiazide diuretics are used to decrease urinary calcium excretion,[5] while antibiotics are administered.

Conventional medications used in the management of kidney stones, including citrate supplements and other pharmacological agents, can be associated with several adverse effects such as nausea, vomiting, diarrhea, and abdominal discomfort. Prolonged use of these medications may also lead to complications, including



hyperkalemia (elevated potassium levels) in patients receiving citrate therapy and metabolic acidosis associated with certain carbonic anhydrase inhibitors. These limitations have encouraged the exploration of alternative therapeutic approaches for the prevention and treatment of urolithiasis.

Herbal medicines have gained considerable attention as potential alternatives to synthetic drugs due to their relatively lower cost, wider availability, and perceived safety profile. Many medicinal plants contain a variety of bioactive compounds that can act on multiple stages of kidney stone formation. These phytochemicals may inhibit crystal nucleation, growth, and aggregation while also providing antioxidant and anti-inflammatory effects that help protect renal tissues.

Among the important phytochemicals reported to possess anti-urolithiatic activity are flavonoids,

polyphenols, tannins, and resveratrol. These compounds have been shown to reduce calcium oxalate crystal formation, minimize oxidative stress, and support normal kidney function. In addition, they exhibit diuretic, antioxidant, and anti-inflammatory properties, which contribute to the prevention and management of urinary stone disease.

Vitis vinifera (grapevine) leaves are known to contain many of these beneficial phytoconstituents. Therefore, the present study was undertaken to evaluate the anti-urolithiatic activity of *Vitis vinifera* leaf extract and to investigate its potential role in the prevention and management of kidney stone formation.

METHODS:

In-Vivo Assay of Anti Urolithiatic Activity of *Vitis vinifera*:

Table No.1: In- Vivo Methods

Sr.No	Test Name/Model	Animal	Principle
1	Ethylene Glycol (EG)– Induced Urolithiasis Model	Wistar rats / Sprague– Dawley rats	EG (0.75–1% in drinking Water) induces hyperoxaluria, causing calcium oxalate stone formation. Test drug prevents or reduces crystal deposition, improves urine/serum markers.[24]
2	Ethylene Glycol + Ammonium Chloride (EG + AC) Model	Wistar rats	AC causes metabolic acidosis and enhances stone formation with EG. Test drug reduces biochemical abnormalities and kidney CaOx deposits.[25]
3	Sodium Oxalate– Induced Urolithiasis	Wistar rats / Mice	Sodium oxalate injection rapidly increases urinary oxalate, causing acute CaOx crystal deposition. Test drug decreases crystalluria and Renal damage.[26]
4	Zinc Disc Implantation Method	Rats	A zinc disc is surgically implanted into the bladder, acting as a nidus for stone growth. Test drug reduces stone size, weight, and deposition rate.[27]

5	Ammonium Oxalate and Vitamin D3–Induced Model	Rats	Vitamin D3 increases calcium absorption; ammonium oxalate increases oxalate load→ CaOx crystals form. Test drug prevents hypercalciuria, stone.[28]
---	---	------	---

In vitro Assay of Anti-Urolithiatic Activity of *Vitis vinifera*:

Table No.2: In-Vitro Methods

Sr. No.	METHODS	PRINCIPLE
1	Aggregation Assay	Induced Model Ethylene glycol causes hyperoxaluria, leading to calcium oxalate stone formation. Test drug prevents or reduces crystal deposition.[29]
2	Dissolution/ Litholytic Assay	Ammonium chloride causes metabolic acidosis, which enhances stone formation with EG. Test drug reduces biochemical abnormalities and CaOx deposits. [30]
3	Microscopic Evaluation	Sodium oxalate injection increases urinary oxalate, rapidly forming CaOx crystals. Test drug decreases crystalluria and kidney damage. [31]
4	Crystal Growth Assay	Assesses whether a test sample inhibits the enlargement of calcium oxalate crystals. [32]
5	Nucleation Assay	Determines how effectively a sample prevents the initial formation of calcium oxalate crystal nuclei. [33]
6	Egg membrane assay	The egg membrane acts as a natural semi-permeable membrane in which calcium oxalate crystals are enclosed. The test drug diffuses through the membrane and dissolves the calcium oxalate crystals. The extent of dissolution indicates anti-urolithiatic activity.

PLANT PROFILE:

GRAPES LEAVES:

Plant description:

Vitis vinifera is a woody climbing plant belonging to the family Vitaceae, commonly known as the grapevine the leaves contain various phytoconstituents such as flavonoids, tannins, polyphenols, saponins, and resveratrol, which are responsible for antioxidant, anti-inflammatory





Fig No. 1: *Vitis vinifera* Leaves

Synonym: European grape vine, Grapevine, wine grape

Scientific name: *Vitis vinifera*

Biological source: *Vitis vinifera* consists of the dried leaves, fruits, and seeds of *Vitis vinifera* Linn, belonging to the family Vitaceae.

Geographical classification: *Vitis vinifera* is native to the Mediterranean region and Western Asia. It is widely cultivated in many countries including India, Italy, France, Spain, the United States, and China. In India, grape cultivation is mainly carried out in Maharashtra, Karnataka, Andhra Pradesh, Tamil Nadu, and Punjab due to favourable climatic conditions.

Scientific classification:

Kingdom	Plantae
Division	Magnoliophyte
Class	Magnoliopsida
Order	Vitales
Family	Vitaceae
Genus	Vitis
Species	Vitis vinifera

Chemical constituents: *Vitis vinifera* contains various important phytochemical constituents responsible for its medicinal properties. The major chemical constituents present in the Leaves, Fruits, and Seeds include Flavonoids, Polyphenols, Tannins, Saponins, Anthocyanins, Terpenoids, Organic Acids, Vitamins, Minerals, and Resveratrol.

Pharmacological Action: Acts as an antioxidant by scavenging free radicals, exhibits anti-

inflammatory activity by reducing inflammatory mediators, shows cardio protective effect by improving blood circulation and reducing oxidative damage and possesses antimicrobial activity against various microorganisms.

PLANT MATERIAL PROCESSING:

Collection of Plant and Authentication:

The leaves samples of the *Vitis vinifera* collected from the nursery and authenticated by Department of Botany, Jaysingpur College, Jaysingpur having



authentication Outward No. AES/JCJ/ By hand. [**Preparation of the plant extract:**
Ref. Annexure No. 1]




Fig No.2: Soxhlet Apparatus



Fig No.3: Extract of *Vitis vinifera*

Yield Calculation:

$$\% \text{Yield} = \text{Weight of Extract (g)} / \text{Weight of dry Powder (g)} \times 100$$

Preliminary Phytochemical Investigation:

Chemical Test:

Various chemical tests are performed for the determination of phytoconstituents present in extract [35]

1) TEST FOR ALKALOIDS:

a) Drangendorff Test:

Take a small quantity of the plant extract in a test tube and dissolve it in a few millilitres of dilute hydrochloric acid. Filter the solution if necessary to obtain a clear extract. Add 2–3 drops of Dragendorff's reagent to the solution and mix gently. Observe the formation of an orange, reddish-brown, or brick-red precipitate. The appearance of this precipitate indicates the presence of alkaloids in the plant extract.

b) Mayer's test:

Take a small amount of the plant extract in a test tube and add a few ml of dilute hydrochloric acid.

Filter the solution if required to obtain a clear extract. Then add 2–3 drops of Mayer's reagent to the solution and shake gently. The formation of a cream or white coloured precipitate indicates the presence of alkaloids in the plant extract.

c) Wagner's test:

Take a small quantity of the plant extract in a test tube and dissolve it in a few ml of dilute hydrochloric acid. Filter the solution if necessary to obtain a clear extract. Add 2–3 drops of Wagner's reagent to the extract and shake gently. The formation of a brown or reddish-brown precipitate indicates the presence of alkaloids in the plant extract.

2) TEST FOR FLAVONOIDS:

a) Lead Acetate Test:

Add a few drops of lead acetate solution to the plant extract. Formation of a yellow precipitate indicates the presence of flavonoids.

b) Pew's Test:

Take a small quantity of the plant extract in a test tube and add a small amount of zinc dust. Then add a few drops of concentrated hydrochloric acid carefully and mix gently. The formation of a red or orange-red colour indicates the presence of flavonoids in the plant extract.

c) Ammonia Test:

Take a small quantity of the plant extract in a test tube and add a few ml of dilute ammonia solution. Then add concentrated sulfuric acid carefully along the sides of the test tube. The appearance of a yellow colour, which disappears on standing, indicates the presence of flavonoids in the plant extract.

3) TANNINS TEST:

a) Gelatine Test:

Take a small quantity of the plant extract in a test tube and add 1% gelatin solution containing sodium chloride. Mix the solution gently and observe the formation of a white precipitate. The appearance of the precipitate indicates the presence of tannins in the plant extract.

b) Braymer's Test:

Take a small quantity of the plant extract in a test tube and add a few drops of 10% alcoholic ferric chloride solution. Mix the contents gently and observe the colour change. The formation of a blue, green, or black coloration indicates the presence of tannins in the plant extract.

c) Bromine Water Test:

Take a small quantity of the plant extract in a test tube and add a few drops of bromine water solution slowly. Mix the solution gently and observe the reaction. The formation of a buff-coloured

precipitate indicates the presence of tannins in the plant Extract.

d) Phenazone Test:

Take a small quantity of the plant extract in a test tube and add sodium acid phosphate solution followed by a few drops of phenazone solution. Mix the contents gently and observe the formation of a bulky precipitate. The appearance of the precipitate indicates the presence of tannins in the plant extract.

4) CARDIAC GLYCOSIDE:

a) Test for Cardenolides:

Pyridine and sodium nitroprusside or sodium NaOH added to extract to observe red or faded to brownish colour.

b) Baljet Test:

Add sodium picrate solution to the plant extract and mix gently. Development of an orange or yellow colour confirms the presence of cardiac glycosides.

c) Keller-Killani Test:

Take a small quantity of the plant extract in a test tube and add glacial acetic acid containing a trace of ferric chloride solution. Carefully add concentrated sulfuric acid along the sides of the test tube. The formation of a brown ring at the junction of the two layers and a bluish-green colour in the upper layer indicates the presence of cardiac glycosides.

5) CARBOHYDRATES:

a) Barfoed's Test:

Add Barfoed's reagent to the plant extract and heat in a water bath. Formation of a red precipitate



within a few minutes indicates the presence of monosaccharides.

b) Seliwanoff's Test:

Take a small quantity of the plant extract in a test tube and add Seliwanoff's reagent. Heat the mixture gently in a water bath for a few minutes. The appearance of a cherry-red colour indicates the presence of ketose sugars such as fructose in the plant extract.

c) Resorcinol Test:

Take a small quantity of the plant extract in a test tube and add a few drops of resorcinol reagent along with concentrated hydrochloric acid. Heat the mixture gently in a water bath for a few minutes. The formation of a red or cherry-red colour indicates the presence of ketose sugars in the plant extract.

IN- VITRO METHODS

Egg Membrane Assay [34]:

Preparation of calcium oxalate stones:

The following procedure was used to make the calcium oxalate (CaOx) experimental kidney stones by homogenous precipitation. In 100 mL of distilled water, 1.47 g of calcium chloride dihydrate was dissolved, and in 100 mL of 1 M H₂SO₄, 1.34 g of sodium oxalate was dissolved. Both were combined in a beaker and stirred to precipitate calcium oxalate. Ammonia solution was used to free the crystals from H₂SO₄. Finally, the crystal was washed in distilled water and dried for four hours at 60°C. Further research was carried out using the artificial urine and stone that had been prepared.




Fig No.4: Calcium Oxalate Crystal

Preparation of the semipermeable membrane from eggs:

The semipermeable membrane was prepared from egg. A glass rod was used to puncture the apex of the eggs and the entire contents were squeezed out. Empty egg shells were washed thoroughly in distilled water, after that the shells were placed in

a beaker containing 4.0 mL concentrated HCl in 200 mL distilled water. The semi-permeable membrane was completely decalcified after being kept for an overnight period. The next day the semi permeable membranes were carefully removed from the egg shell and washed thoroughly with distilled water. The trace of acid present in the membrane is neutralised by placing the shell

membranes in an ammonia solution, and rinsed with distilled water. It was kept in a moistened state in the refrigerator at a pH of 7–7.4.



Fig No.5: Decalcification of Egg Membrane



Fig No.6: Semipermeable Membrane of Eggs

Methods:

The studies were carried out in three groups as per experimental design

- Group 1: Negative Control (1 mg Calcium Oxalate)
- Group 2: 1 mg Calcium Oxalate +10 mg of Standard (Cystone)
- Group 3: 1 mg Calcium Oxalate +10 mg of Leaves Extract

All the above groups were packed in semi permeable membrane by suturing. They were suspended in a conical flask containing 100 ml of 0.1 M TRIS buffer. The conical flasks of all groups were placed in an incubator and preheated to 37°C for 2 hours. The contents of semi permeable membrane from each group were taken into test tubes. 2 ml of sulphuric acid was added to each test tube and titrated with 0.2 M KMnO_4 till a light pink colour end point was obtained. 1 ml of 0.2 M KMnO_4 is equivalent to 0.1898 mg of calcium. Percentage dissolution of calcium oxalate in various groups was calculated.




Fig No. 7: In –Vitro model used to evaluate anti-urolithiatic activity

Crystal Nucleation Assay:

A buffer containing 0.05 M Tris–HCl and 0.15 M NaCl at pH 6.5, a solution of calcium chloride (5 mM) and sodium oxalate (7.5 M) was prepared. A 9 mL calcium chloride solution was combined with 1 mL extracts at various concentrations (50, 100,250, 500, and 1000 mg/mL). Crystallization was stimulated by the insertion of 1 ml of sodium oxalate solution and the absorbance shift was recorded at 620 nm in a UV spectrophotometer

(UV- 1800, Shimadzu Pvt. Ltd.) for 30 minutes at 37oC.

The procedure was followed for the control, substituting distilled water instead of the extract. All samples were inspected in triplicate. Standard drug Cystone was used as a positive control for comparison at distinct concentrations including 50, 100,250, 500, and 1000 mg/mL. The percentage inhibition of nucleation rate was calculated by given formula

$$\% \text{ Inhibition} = \frac{\text{Absorbance of Control} - \text{Absorbance of Test}}{\text{Absorbance of Control}} \times 100$$

RESULTS:

Collection and Authentication of plant material:

The whole plant was identified and authenticated by Prof. Manisha V. Kale Head Dept. of Botany Jaysingpur College, Jaysingpur.

Preparation of Plant Extract:

The total yield of Ethanolic Extract was found to be

Table No.3: % yield of Ethnolic Extract

Sr. No	Extract	% Yield (w/w)
1.	Leaves	2.7

PRELIMINARY PHYTOCHEMICAL INVESTIGATION:

Table No.4: Phytochemical evaluation test:

Sr. No.	Test	EEVV
1	Test for Alkaloids	
	Mayer's test	+
	Wagner's test	+
	Dragendroff's test	+
	Hager's test	-
2	Test for flavonoids	
	Lead acetate test	+
	Pew's test	+
	Ammonia test	+
	Ferric chloride test	-
3	Test for Glycosides	
	Cardenoloides test	+
	Baljet test	+
	Keller killani test	+
	Kedee's test	-
4	Test for Carbohydrates	
	Barfoed test	+
	Seliwanoff's test	+
	Resorcinol test	+
	Test for pentose	-
5	Test for saponins	
	Foam test	-
6	Test for tannins	
	Gelatine test	+
	Braymer's test	+
	Bromine water test	+
	Phenadzone test	+





Fig. No.8: Alkaloid Test




Fig. No.9: Flavonoids test




Fig. No.10: Glycosides Test



Fig. No. 11: Carbohydrates Test




Fig. No. 12: Tannins Test

Various chemical tests are performed for the phytochemical investigation. From the results as we found that all chemical tests are positive for the flavonoids, alkaloids, tannins, carbohydrate test in *Vitis vinifera* leaves Which indicates that the major phyto-constituents like flavonoids, alkaloids, carbohydrate, glycoside and tannins are present in the leaves extract of *Vitis vinifera*.

In-Vitro Evaluation of Anti-Urolithiatic Activity

Egg Membrane Assay

Groups as per experimental design

- Group 1: Negative Control (1 mg Calcium Oxalate)
- Group 2: 1 mg Calcium Oxalate +10 mg of Standard (Cystone)
- Group 3: 1 mg Calcium Oxalate +10 mg of Leaves Extract

Table No.5: Percent of Calcium Oxalate dissolution by various groups under study

Groups	Vol. of Standard KMnO4(ml)	Wt. Of Calcium Oxalate Taken	Wt. Of Calcium Estimated (Mg)	Wt. Of Calcium Reduced (Mg)	% Dissolution
1	0.1	1	-	-	-
2	0.6	1	0.11388	0.88612	88.61
3	0.9	1	0.17082	0.82918	82.91




Fig. No. 13: Percentage Dissolution of Calcium Oxalate by Various Groups under Study

The egg membrane assay showed considerable dissolution of calcium oxalate crystals in the treated groups. The standard drug (Cystone)

exhibited 88.61% dissolution, while the Ethanolic extract of *Vitis vinifera* leaves showed 82.91% dissolution of calcium oxalate crystals. The



findings indicate that the leaves extract possesses **Nucleation Assay:** significant Anti-urolithiatic Activity.

Table N0.6: Nucleation Assay of Standard (Cystone) drug

GROUP	CONCENTRATION $\mu\text{g/mL}$	ABSORBANCE	%INHIBITION	IC 50
Standard (Cystone)	50	0.486 \pm 0.0004	48.44	56.5 $\mu\text{g/}$ mL
	100	0.373 \pm 0.0008	60.48	
	250	0.297 \pm 0.0008	68.58	
	500	0.177 \pm 0.0004	81.17	
	1000	0.107 \pm 0.0012	88.59	




Fig. No. 14: Cystone (Standard Drug) at Various Concentration

Table No.7: Nucleation Assay of Ethanolic extract of *Vitis vinifera* leaves

GROUP	CONCENTRATION $\mu\text{g/mL}$	ABSORBANCE	%INHIBITION	IC50
TEST (EEVV)	50	0.566 \pm 0.0001	39.97	78.6 $\mu\text{g/m}$ L
	100	0.401 \pm 0.0004	57.48	
	250	0.375 \pm 0.0008	60.27	
	500	0.302 \pm 0.0012	67.93	
	1000	0.217 \pm 0.001	76.97	

The results of the nucleation assay demonstrated that the standard drug Cystone exhibited significant anti-urolithiatic activity by inhibiting calcium oxalate crystal nucleation in a concentration-dependent manner. The percentage

inhibition is 48.44, 60.48, 68.58, 81.17, 88.59 at concentration 50, 100, 250, 500, 1000 $\mu\text{g/mL}$ respectively the IC₅₀ value of 56.5 $\mu\text{g/mL}$ suggests good inhibitory potential against kidney stone formation






Fig No. 15: Ethanolic extract of *Vitis vinifera* leaves at various concentration

DISCUSSION:

Kidney stone formation mainly occurs due to supersaturation and crystallization of calcium oxalate in the urinary tract. Therefore, inhibition of calcium oxalate crystal formation is considered an important approach in the prevention and treatment of urolithiasis. The present study was carried out to evaluate the anti-urolithiatic activity of the ethanolic extract of *Vitis vinifera* leaves using different in-vitro models such as phytochemical screening, egg membrane assay, and crystal nucleation assay

The preliminary phytochemical investigation of the ethanolic extract revealed the presence of important phytoconstituents such as flavonoids, alkaloids, tannins, glycosides, and carbohydrates. These phytochemicals are known to possess antioxidant, anti-inflammatory, and diuretic properties which may contribute to anti-urolithiatic activity. Flavonoids and polyphenolic compounds such as catechin and Gallic acid help reduce oxidative stress and prevent crystal aggregation, while tannins such as Hydrolysable

tannins may inhibit calcium oxalate crystal formation.

In the egg membrane assay, the ethanolic extract showed significant dissolution of calcium oxalate crystals. The extract produced 82.91% dissolution, whereas the standard drug Cystone showed 88.61% dissolution. The results indicate that the extract possesses considerable litholytic activity and can dissolve preformed calcium oxalate crystals effectively. Although the activity was slightly lower than the standard drug, the extract demonstrated promising anti-urolithiatic potential.

The nucleation assay further confirmed the anti-urolithiatic activity of the extract. The ethanolic extract inhibited calcium oxalate crystal nucleation in a concentration-dependent manner. The percentage inhibition increased from 39.97% at 50 µg/mL to 76.97% at 1000 µg/mL. Similarly, the standard drug Cystone showed inhibition ranging from 48.44% to 88.59% at corresponding concentrations.

The IC₅₀ value of the ethanolic extract was found to be 78.6 µg/mL, while the standard drug showed an IC₅₀ value of 56.5 µg/mL. A lower IC₅₀ value of



the standard indicates stronger activity; however, the extract still exhibited good inhibitory potential against calcium oxalate crystallization. The activity of the extract may be attributed to the synergistic action of flavonoids, tannins, saponins, and other phenolic compounds present in the leaves.

Overall, the findings of the study suggest that the ethanolic extract of *Vitis vinifera* leaves possesses significant anti-urolithiatic activity by inhibiting nucleation and promoting dissolution of calcium oxalate crystals. The study supports the traditional use of herbal medicines in the management of kidney stones and indicates that the plant may serve as a potential natural therapeutic agent for urolithiasis. However, further in-vivo studies and clinical investigations are required to confirm its safety, efficacy, and mechanism of action.

CONCLUSION:

The present study demonstrated that the ethanolic extract of *Vitis vinifera* leaves possesses significant anti-urolithiatic activity against calcium oxalate crystal formation. Phytochemical screening confirmed the presence of important bioactive constituents such as flavonoids, tannins, alkaloids, glycosides, and carbohydrates, which may contribute to its therapeutic effect. In the in-vitro evaluation, the extract showed considerable calcium oxalate crystal dissolution in the egg membrane assay and effectively inhibited crystal nucleation in a concentration-dependent manner, although slightly lower than the standard drug Cystone.

The findings suggest that *Vitis vinifera* leaves have promising potential as a natural remedy for the prevention and management of kidney stones due to their crystal inhibitory and antioxidant properties. However, further in-vivo and clinical studies are required to establish their safety,

efficacy, and mechanism of action for therapeutic use.

REFERENCES

1. Soriano RM, Penfold D, Leslie SW. Anatomy, abdomen and pelvis, kidneys.
2. Allam EA. Urolithiasis unveiled: pathophysiology, stone dynamics, types, and inhibitory mechanisms: a review. African Journal of Urology. 2024 Jul 23;30(1):34
3. rights are reserved by PRADNYA A, JAGTAP N. Comprehensive Review on Kidney Stone.
4. Khan F, Haider MF, Singh MK, Sharma P, Kumar T, Neda EN. A comprehensive review of kidney stones, its diagnosis and treatment with allopathic and ayurvedic medicines. Urol Nephrol Open Access J. 2019 Aug;7(4):69-74.
5. Holmes RP, Goodman HO, Assimos DG. Contribution of dietary oxalate to urinary oxalate excretion. Kidney Int. 2001;59(1):270-276.
6. Barbasa C, Garciaa A, Saavedra L, et al. Urinary analysis of nephrolithiasis markers. J Chromatogr B 2002;781:433-55.
7. Arya P, Pandey S, Verma V. Kidney stone formation and use of medicinal plants as antiurolithiatic agents. Universal Journal of Pharmaceutical Research 2017;
8. McAteer JA, Evan AP, Williams JC Jr, Lingeman JE. Treatment protocols to reduce renal injury during shock wave lithotripsy. Curr Opin Urol 2009 Mar; 19(2): 192-5. PMID: 19195131
9. <http://www.imop.gr/en/uoinfo-urolithiasis%23#antimetopisi>
10. APA. Tripathi, K. D. (2018). Essentials of medical pharmacology (8th ed.)
11. Abdel-Salam NA, Ghazy NM, Sallam SM, Radwan MM, Wanas AS, ElSohly MA, El-L. and their immune stimulant, antioxidant and cytotoxic activities on hepatocellular



- carcinoma HepG-2 cell line. Natural product research. 2018 Mar 19;32(6):702-6.
12. Akram M, Asif HM, Akhtar N, Shah PA, Uzair MU, Shaheen G, Shamim T, Shah SA, Ahmad K. *Tribulus terrestris* Linn.: a review article. *J Med Plants Res.* 2011 Aug 18;5(16):3601-5.
 13. Elufioye TO, Oyediji AO, Habtemariam S. A review of the traditional uses, phytochemistry and pharmacology of *Bryophyllum pinnatum* (Lam.)(Crassulaceae). *Journal of Biologically Active Products from Nature.* 2022 May 4;12(3):190-222.
 14. Kumar K, Raza MA, Yadav N, Chaurasiya J, Dwivedi KP. A Brief Study on Fenugreek (*Trigonella Foenum-Graecum*): A.
 15. Salama, Mahitab I. El-kassaby, Azza Hassan Published in *Egyptian Pharmaceutical...* 1 October 2019 *Medicine, Environmental Science*
 16. Kanu CL, Owoeye O, Imosemi IO, Malomo AO. A review of the multifaceted usefulness of *Celosia argentea* Linn. *European Journal of Pharmaceutical and Medical Research.* 2017;4(10):72-9.
 17. Debashri M, Tamal M. A Review on efficacy of *Azadirachta indica* A. Juss based biopesticides: An Indian perspective. *Res. J. Recent Sci.* 2012;2277:2502.
 18. Bhokardankar PS, Mane SG, Khairanar BP. An overview of shatavari (*asparagus racemosus*) an ayurvedic drug. *International Journal of Ayurveda and Pharma Research.* 2019 Sep 25:60- 5.
 19. Gour R. *Boerhaavia diffusa* Linn plant: a review plant with many therapeutic uses. *Int. J Pharm. Sci. Med.* 2021;6(4):25-41.
 20. Bolaji AO, Oladejo AS, Adeniran OI. Morphological characterization and cytological studies of the green-stemmed and the red-stemmed *Basella alba*. *Notulae Scientia Biologicae.* 2023 Jun 19;15(2):11338
 21. Hardegger M, Sturm A. Transformation and regeneration of carrot (*Daucus carota* L.). *Molecular Breeding.* 1998 Apr;4(2):119-27.
 22. Nimavat A, Trivedi A, Yadav A, Patel P. A review on kidney stone and its herbal treatment. *Journal of Pharmacy and Pharmacology.* 2022;10:195-209.
 23. Kumar P, Singh K, Gairola S. Botanical standardization of raw herbal drug *Pashanabheda* [*Bergenia ciliata* (Haw.) Sternb.] used in Indian systems of medicine.
 24. Prasad KVS et al. Evaluation of antiurolithiatic activity of *Tribulus terrestris* in EG-induced urolithiasis in rats. *J Ethnopharmacol.* 2007.
 25. Haritha C et al. EG + AC-induced hyperoxaluria model for screening antiurolithiatic drugs. *Indian J Pharm Sci.* 2014.
 26. Lakshmi V et al. Sodium oxalate-induced urolithiasis model in rats. *Pharmacologyonline* 2011 Atmani F et al. Zinc disc implantation model for bladder stone formation. *Urol Res.* 2004.
 27. Selvam R. Prevention of vitamin D3 + ammonium oxalate-induced urolithiasis by natural extracts. *Phytother Res.* 2001.
 28. Divakar K et al. Gentamicin-induced renal calculi model. *J Biol Sci.* 2010. Alelign T, Petros B. Kidney stone disease: an update on current concepts. *Advances in Urology.* 2018;2018:3068365.
 29. IJLPR Journal, "In vitro evaluation of antiurolithiatic activity of *Asparagus racemosus*", 2019
 30. *Journal of Pharma Sci Innova*, "Evaluation of in vitro antiurolithiatic activity of *Pavonia zeylanica*", 2020.
 31. *Ceterach officinarum* alters CaOx morphology
 32. *Pharmacognosy Journal*, "Antiurolithiatic activity of *Daucus carota*: An in vitro study", 2018.



33. Saha S, Verma RJ. Inhibition of calcium oxalate crystallization in vitro by an extract of *Bergenia ciliata*. *Arab Journal of Urology*. 2013.
34. <https://www.sciencedirect.com/science/article/pii/S2772371223000463?utm>
35. Pooja Verma A, Lal VK. Punarnava—A natural remedy by Ayurveda. *Int J Pharm Sci*. 2014
36. Kokate CK, Purohit AP, Gokhale SB. *Pharmacognosy*. Nirali Prakashan.

HOW TO CITE: Sumit Shinge, Sumayya Shaikh, Sujit Chavan, Sujit Ukande, Sushant Patil, Evaluation of Anti Uroethiatic Potential of *Vitis Vinifera* Leaves, *Int. J. of Pharm. Sci.*, 2026, Vol 4, Issue 6, 6353-6371. <https://doi.org/10.5281/zenodo.20844672>

