INTERNATIONAL JOURNAL OF PHARMACEUTICAL SCIENCES
[ISSN: 0975-4725; CODEN(USA): IJPS00]
Journal Homepage: https://www.ijpsjournal.com

OPEN ACCESS

Review Article

Formulation and Evaluation of a Herbal Analgesic Cream Containing Menthol, Camphor, and Clove Oil

Ankita soni*[1], Atul Bisen[2], Rajesh Mujariya[3], Manjeet Singh[4]

[1]*Research Scholar of Institute of Pharmaceutical Science & Research, Balaghat (M.P.)*
[2]*Associate, Professor of Institute of Pharmaceutical Science & Research, Balaghat (M.P.)*
[3]*Principal & Professor of Institute of Pharmaceutical Science & Research, Balaghat (M.P.)*
[4]*Executive Director of Institute of Pharmaceutical Science & Research, Balaghat (M.P.)*

ARTICLE INFO

Published: 11 May. 2026
Keywords:
Herbal analgesic, Menthol, Camphor, Clove oil, Topical cream, Pain management, Phytopharmaceuticals
DOI:
10.5281/zenodo.20126486

ABSTRACT

Pain is a common clinical condition often associated with inflammation, injury, or chronic disorders, significantly impacting quality of life. Conventional analgesics such as NSAIDs are effective but frequently produce adverse effects during long-term use, including gastrointestinal and renal complications. This has increased interest in safer, plant-based alternatives, particularly topical herbal formulations that provide localized action with minimal systemic exposure. Menthol, camphor, and clove oil are widely recognized phytoconstituents with significant analgesic and anti-inflammatory properties. Menthol induces a cooling sensation by activating TRPM8 receptors, camphor acts as a counter-irritant stimulating sensory nerve endings, and clove oil, rich in eugenol, exhibits analgesic effects through prostaglandin inhibition and local anesthetic action. The development of an effective herbal analgesic cream requires appropriate selection of excipients, emulsification techniques, and optimization of physicochemical parameters such as pH, viscosity, and spreadability. Evaluation involves physical characterization, stability studies, and in vitro and in vivo assessments to ensure safety and efficacy.

INTRODUCTION

1.1 Pain and Inflammation

Pain is an unpleasant sensory and emotional experience associated with actual or potential tissue damage. It serves as a protective mechanism, alerting the body to injury or disease. Based on duration and underlying cause, pain is broadly classified into acute pain, which is short-term and usually linked to tissue injury; chronic pain, which persists beyond normal healing time

***Corresponding Author:** Ankita soni
Address: *Research Scholar of Institute of Pharmaceutical Science & Research, Balaghat (M.P.)*
Email ✉: ankitasoni2803@gmail.com
Relevant conflicts of interest/financial disclosures: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

and may last for months or years; and neuropathic pain, which arises from damage or dysfunction of the nervous system.[1] Inflammation is a complex biological response of body tissues to harmful stimuli such as pathogens, damaged cells, or irritants. It is characterized by redness, swelling, heat, pain, and loss of function. [2] The pathophysiology of pain and inflammation involves the release of various chemical mediators, including prostaglandins, cytokines, histamine, and bradykinin. Prostaglandins, synthesized via the cyclooxygenase (COX) pathway, play a crucial role in sensitizing nociceptors and enhancing pain perception. Similarly, pro-inflammatory cytokines such as interleukins and tumor necrosis factor (TNF-α) contribute to the initiation and maintenance of inflammatory responses, thereby intensifying pain. [3]

1.2 Limitations of Conventional Analgesics

Conventional analgesics, particularly non-steroidal anti-inflammatory drugs (NSAIDs), are widely used for the management of pain and inflammation. [4] These drugs exert their effect primarily by inhibiting COX enzymes and reducing prostaglandin synthesis. However, prolonged use of NSAIDs is associated with several adverse effects, including gastrointestinal irritation, ulceration, bleeding, and renal toxicity. In some cases, they may also increase the risk of cardiovascular complications. [5] These limitations highlight the need for safer and more effective alternatives that can provide adequate pain relief without causing significant side effects. The growing awareness regarding drug-related toxicity has shifted attention toward natural and plant-based therapies. [6]

1.3 Role of Herbal Medicine in Pain Management

Herbal medicine has been used for centuries in traditional systems such as Ayurveda, Unani, and Traditional Chinese Medicine for the treatment of pain and inflammation. [7] In recent years, there has been a significant increase in the global demand for herbal formulations due to their perceived safety, efficacy, and natural origin. Herbal drugs offer several advantages, including lower incidence of side effects, cost-effectiveness, and better patient compliance. Many medicinal plants contain bioactive compounds such as flavonoids, alkaloids, terpenoids, and phenolics, which exhibit analgesic and anti-inflammatory properties. These compounds often act through multiple mechanisms, providing a holistic approach to pain management. [8]

1. Rationale of Polyherbal Analgesic Cream

The concept of polyherbal formulation is based on the principle of synergistic action, where the combined effect of multiple herbal ingredients is greater than the sum of their individual effects. In a polyherbal analgesic cream containing menthol, camphor, and clove oil, each component contributes through a distinct mechanism, resulting in enhanced therapeutic efficacy. [9] Topical drug delivery offers several advantages over oral administration. It avoids first-pass metabolism, thereby improving drug availability at the site of action. It provides localized action, directly targeting the affected area, which leads to faster relief from pain and inflammation. Additionally, topical formulations reduce systemic side effects and improve patient compliance, making them a suitable choice for long-term use. [10]

2. Plant Profile

2.1 Menthol

Menthol is a naturally occurring compound obtained from peppermint oil and is widely used in topical formulations due to its cooling and soothing properties. When applied to the skin, it produces a characteristic cooling sensation that helps in relieving pain and discomfort. [11]

Pharmacologically, menthol acts by stimulating cold-sensitive receptors in the skin, particularly TRPM8 channels, which create a cooling effect and reduce the perception of pain. [12] It also exhibits mild anti-inflammatory activity and acts as a counter-irritant, diverting attention from deeper pain. Due to these properties, menthol is commonly used in analgesic creams, gels, and balms for the management of muscle pain, joint pain, and minor injuries. [13]

Figure no. 1: Menthol

2.2 Camphor [14]

Camphor is a terpenoid compound traditionally obtained from the wood of *Cinnamomum camphora*. It is widely used in topical medicinal preparations because of its analgesic and anti-inflammatory properties. Camphor acts mainly as a counter-irritant by stimulating sensory nerve endings, which helps in reducing pain perception. It produces a warming or cooling sensation depending on its concentration. Additionally, camphor improves local blood circulation, which may contribute to faster relief from pain and inflammation. However, it should be used within safe limits, as higher concentrations may cause skin irritation or toxicity.

Figure no. 2: Camphor

2.3 Clove Oil

Clove oil is extracted from the dried flower buds of *Syzygium aromaticum* and is well known for its strong analgesic and antiseptic properties. [15] The major active constituent of clove oil is eugenol, which is responsible for most of its pharmacological effects. [16] Eugenol exhibits significant analgesic activity by inhibiting the synthesis of prostaglandins and acting as a local anesthetic. [17] It also possesses anti-inflammatory, antioxidant, and antimicrobial properties. Due to these effects, clove oil is widely used in dental preparations and topical formulations for the relief of muscular and joint pain. [18]

Figure no. 3: Clove Oil

3. Mechanism of Analgesic Action

The analgesic effect of a polyherbal cream containing menthol, camphor, and clove oil is achieved through multiple complementary mechanisms. [19] Menthol works by activating TRPM8 receptors, producing a cooling sensation that reduces pain perception. Camphor acts as a counter-irritant, stimulating nerve endings and creating a mild warming or cooling effect, which helps to distract from deeper pain signals. It may also improve local blood circulation. [20] Clove oil, through its active constituent eugenol, inhibits the cyclooxygenase (COX) pathway, thereby reducing the production of prostaglandins, which are key mediators of pain and inflammation. It also provides a local anesthetic effect by numbing the affected area. [21] Together, these ingredients produce a synergistic effect by acting through different pathways, including sensory modulation, counter-irritation, and inhibition of inflammatory mediators. This multi-mechanistic approach enhances the overall analgesic efficacy while minimizing dependence on synthetic drugs. [22]

Table 1: Mechanism of Action of Active Ingredients

Ingredient	Mechanism	Pharmacological Effect
Menthol	TRPM8 receptor activation	Cooling, analgesic
Camphor	Counter-irritant action	Pain relief, improved circulation
Clove Oil (Eugenol)	COX inhibition	Analgesic, anti-inflammatory

4. Formulation Aspects of Herbal Cream

4.1 Types of Cream Base

Creams are semi-solid emulsions that can be classified based on the nature of the continuous phase. The selection of an appropriate base plays a crucial role in determining the stability, absorption, and effectiveness of the formulation. [23]

Oil-in-water (O/W) creams consist of oil droplets dispersed in a continuous aqueous phase. These creams are non-greasy, easily washable, and provide a cooling effect on the skin. Due to their better patient acceptability and ease of application, O/W creams are commonly preferred for topical analgesic formulations.

Water-in-oil (W/O) creams contain water droplets dispersed in an oil phase. These creams are more occlusive and provide better moisturization, making them suitable for dry or scaly skin conditions. However, they tend to be greasy and less comfortable for frequent use compared to O/W creams. [24]

4.2 Excipients Used

Excipients are inactive substances that play a vital role in enhancing the stability, consistency, and performance of the cream formulation.

Emulsifiers, such as stearic acid and cetyl alcohol, help in stabilizing the emulsion by reducing surface tension between oil and water phases. They also contribute to the texture and viscosity of the cream. [25]

Preservatives, including parabens, are added to prevent microbial growth and extend the shelf life of the product, especially since creams contain water which can support microbial contamination.

Humectants, such as glycerin and propylene glycol, help in retaining moisture by attracting water to the skin. They improve skin hydration and enhance the overall feel and spreadability of the formulation.

4.3 Method of Preparation

The preparation of herbal creams involves a systematic process to ensure uniform distribution of ingredients and stability of the final product. In the fusion method, the oil phase (containing lipophilic ingredients) and aqueous phase are heated separately to the same temperature, typically around 70–75°C. The aqueous phase is then slowly added to the oil phase with continuous stirring until a uniform emulsion is formed. [26] The emulsification process involves constant mixing to reduce droplet size and ensure proper dispersion of one phase into the other, resulting in a stable and homogeneous cream. After the base is prepared and cooled to a suitable temperature, the active ingredients such as menthol, camphor, and clove oil are incorporated with gentle stirring to avoid loss of volatile components and to maintain their therapeutic activity. [27]

Figure 4: General Method of Cream Preparation

Oil Phase Preparation (Stearic acid + Cetyl alcohol + Oils)
↓ (Heat 70–75°C)
Water Phase Preparation (Water + Glycerin + Preservatives)
↓ (Heat 70–75°C)
Mixing of Both Phases
↓
Continuous Stirring
↓
Cooling of Emulsion
↓
Addition of Menthol + Camphor + Clove Oil
↓
Final Herbal Cream

5. Evaluation Parameters of Cream

5.1 Physical Evaluation

Physical evaluation is essential to assess the appearance and acceptability of the cream formulation. The appearance, including color and odor, should be uniform and pleasant without any signs of phase separation. The pH of the cream should be compatible with skin (generally between 5.5 and 7) to avoid irritation. Homogeneity is checked by visual inspection to ensure uniform distribution of all ingredients without lumps or grittiness. [28]

5.2 Rheological Properties Rheological evaluation helps in determining the flow behavior and application characteristics of the cream.

Viscosity is an important parameter that affects the consistency and stability of the formulation. It is usually measured using a viscometer. [29]

Spreadability indicates how easily the cream can be applied on the skin. A good formulation should spread smoothly with minimal effort, ensuring better patient compliance and uniform application. [30]

5.3 Stability Studies

Stability studies are carried out to evaluate the physical and chemical stability of the formulation over time.

Accelerated stability testing involves storing the cream at different temperature and humidity

conditions to predict its shelf life and stability under normal conditions.

Phase separation is monitored during storage to ensure that the emulsion remains stable and does not break into separate oil and water phases. The absence of phase separation indicates good formulation stability. [31]

Table 2: Evaluation Parameters of Cream

Parameter	Method/Instrument	Ideal Result
Appearance	Visual inspection	Smooth, uniform
Color & Odor	Organoleptic evaluation	Pleasant
pH	Digital pH meter	5.5 – 7
Homogeneity	Visual/Touch	No lumps
Viscosity	Brookfield viscometer	Moderate
Spreadability	Glass slide method	Good
Stability	Accelerated conditions	No phase separation

6. Advantages of Herbal Analgesic Cream

Herbal analgesic creams offer several benefits over conventional synthetic formulations, making them increasingly popular in pain management. [32] One of the major advantages is the reduced risk of side effects, as herbal ingredients are generally better tolerated by the skin and body. This makes them suitable for long-term use, especially in chronic conditions. [33] Another important benefit is better patient compliance. Due to their natural origin, pleasant aroma, and soothing effect, patients are more willing to use herbal formulations regularly. [34] Additionally, such creams often exhibit multifunctional activity. Along with providing analgesic effects, they may also possess antimicrobial, anti-inflammatory, and antioxidant properties. This combined action not only helps in pain relief but also supports overall skin health and healing. [35]

7. Challenges and Limitations

Despite their advantages, herbal analgesic creams also face certain challenges. One of the major issues is the standardization of herbal ingredients, as the composition of plant extracts can vary depending on factors such as geographical source, harvesting conditions, and processing methods. Stability is another concern, particularly due to the presence of volatile oils like menthol, camphor, and clove oil, which may evaporate or degrade over time, affecting the efficacy of the formulation. There is also a possibility of skin irritation or sensitization, especially when used in higher concentrations or in sensitive individuals. Therefore, proper formulation and safety evaluation are essential. Furthermore, the lack of well-designed clinical trials limits the scientific validation and wider acceptance of these formulations in mainstream medicine.

8. Regulatory Aspects

Herbal formulations in India are regulated under guidelines provided by the Ministry of AYUSH. These guidelines ensure the safety, quality, and efficacy of herbal products through proper documentation and standardization. In addition, phytopharmaceutical regulations have been introduced to provide a more structured framework for the development and approval of plant-based drugs. These regulations emphasize scientific evidence, including pharmacological, toxicological, and clinical data.

Quality control is a critical component of regulatory compliance. It involves testing for

parameters such as purity, identity, strength, and microbial contamination to ensure consistency and safety of the final product.

9. Future Perspectives

The future of herbal analgesic creams lies in the integration of advanced drug delivery systems. Development of nanoemulsions and nano-creams can improve the stability, penetration, and bioavailability of active ingredients. Controlled drug delivery systems may further enhance therapeutic efficacy by providing sustained release of active compounds over an extended period. There is also a strong need for clinical validation studies to establish the safety and effectiveness of these formulations in human subjects. Such studies will help in gaining wider acceptance in the medical community. With increasing demand for natural products, there is significant potential for commercial scalability, provided that issues related to standardization and quality control are effectively addressed.

CONCLUSION

Herbal analgesic creams containing menthol, camphor, and clove oil represent a promising alternative for the management of pain and inflammation. Their ability to act through multiple mechanisms provides effective relief while minimizing the side effects commonly associated with synthetic drugs. The concept of polyherbal formulation enhances therapeutic efficacy through synergistic action of different phytoconstituents. At the same time, topical delivery ensures targeted action and improved patient acceptability. However, further research is necessary to overcome existing challenges, particularly in terms of standardization, stability, and clinical validation. With continued scientific advancements, herbal analgesic creams have strong potential to become a reliable and widely accepted option in Modern Pain Management.

REFERENCES

1. Raja SN, Carr DB, Cohen M, Finnerup NB, Flor H, Gibson S, et al. The revised definition of pain. *Pain*. 2020;161(9):1976–82.
2. Ricciotti E, FitzGerald GA. Prostaglandins and inflammation. *Arterioscler Thromb Vasc Biol*. 2011;31(5):986–1000.
3. Medzhitov R. Origin and physiological roles of inflammation. *Nature*. 2008;454(7203):428–35.
4. Vane JR, Botting RM. Mechanism of action of NSAIDs. *Am J Med*. 1998;104(3A):2S–8S.
5. Bjarnason I, Scarpignato C, Holmgren E, et al. NSAIDs and gastrointestinal toxicity. *Gastroenterology*. 2018;154(3):500–14.
6. Grosser T, Smyth E, FitzGerald GA. Anti-inflammatory agents. *Goodman & Gilman's*. 2018.
7. Ekor M. The growing use of herbal medicines. *Front Pharmacol*. 2014;4:177.
8. Yuan H, Ma Q, Ye L, Piao G. Natural products in medicine. *Molecules*. 2016;21(5):559.
9. Fabricant DS, Farnsworth NR. Value of plants in medicine. *Environ Health Perspect*. 2001;109(Suppl 1):69–75.
10. WHO. Traditional medicine strategy 2014–2023. Geneva; 2013.
11. Eccles R. Menthol pharmacology. *J Pharm Pharmacol*. 1994;46(8):618–30.
12. Patel T, Ishiuji Y, Yosipovitch G. Menthol cooling effects. *J Am Acad Dermatol*. 2007;57(5):873–8.
13. Galeotti N, Mannelli LD, Mazzanti G, et al. Menthol analgesic activity. *Neurosci Lett*. 2002;322(3):145–8.
14. Xu H, Blair NT, Clapham DE. Camphor and TRP channels. *J Neurosci*. 2005;25(39):8924–37.

15. Zu Y, Yu H, Liang L, et al. Clove oil pharmacology. *Phytother Res*. 2010;24(4):595–601.
16. Pramod K, Ansari SH, Ali J. Eugenol pharmacology. *Nat Prod Commun*. 2010;5(12):1999–2006.
17. Chaieb K, Hajlaoui H, Zmantar T, et al. Clove oil activity. *Phytother Res*. 2007;21(6):501–6.
18. Marchese A, Barbieri R, Coppo E, et al. antimicrobial activity of eugenol. *Molecules*. 2017;22(5):1–21.
19. Julius D, Basbaum AI. Molecular pain mechanisms. *Nature*. 2001;413(6852):203–10.
20. Caterina MJ, Schumacher MA, Tominaga M, et al. TRPV channels. *Nature*. 1997; 389:816–24.
21. Smith WL, DeWitt DL. Cyclooxygenase pathway. *Annu Rev Biochem*. 2000; 69:145–82.
22. Rang HP, Dale MM, Ritter JM, Flower RJ. *Rang & Dale Pharmacology*. 8th ed. 2016.
23. Aulton ME, Taylor KMG. *Aulton's Pharmaceutics*. Elsevier; 2018.
24. Sinko PJ. *Martin's Physical Pharmacy*. 6th ed. 2011.
25. Lachman L, Lieberman HA. *Industrial Pharmacy*. 3rd ed. 2009.
26. Tadros T. Emulsion formation. Wiley; 2013.
27. Schramm LL. Emulsions and creams. *Adv Colloid Interface Sci*. 2005; 123:231–49.
28. ICH Q1A(R2). Stability testing guidelines. 2003.
29. Allen LV. Dosage forms. 10th ed. 2013.
30. Banker GS, Rhodes CT. Modern pharmaceutics. 2002.
31. Sinko PJ. Rheology in dosage forms. 2011.
32. Mukherjee PK. Herbal drug standardization. Elsevier; 2019.
33. Parasuraman S, Thing GS. Polyherbal concept. *J Tradit Complement Med*. 2014;4(3):143–7.
34. Wagner H. Synergy research. *Fitoterapia*. 2011;82(1):34–7.
35. Williamson EM. Synergy in phytomedicine. *Phytomedicine*. 2001;8(5):401–9.
36. Benson HA. Transdermal drug delivery. *Curr Drug Deliv*. 2005;2(1):23–33.
37. Prausnitz MR, Langer R. Transdermal delivery. *Nat Biotechnol*. 2008;26(11):1261–8.
38. Barry BW. Dermatological formulations. *Drugs Pharm Sci*. 2001.
39. Ekor M. Herbal toxicity issues. *Front Pharmacol*. 2014; 4:177.
40. WHO. GACP guidelines. 2003.
41. Barnes J. Herbal medicine safety. *Pharm J*. 2003.
42. Ministry of AYUSH. Ayurvedic Pharmacopoeia of India.
43. CDSCO. Phytopharmaceutical guidelines India.
44. WHO. Quality control of herbal medicines. 2011.
45. Gupta PK, Pandit JK. Nanotechnology in pharmaceutics. *Int J Pharm*. 2010; 386:1–11.
46. Shakeel F, Ramadan W. Nanoemulsion systems. *AAPS PharmSciTech*. 2010;11(2):674–81.
47. Date AA, Naik B, Nagarsenker MS. Nanoemulsions. *Int J Pharm*. 2010; 355:19–30.
48. Patel VR, Agrawal YK. Nanotechnology applications. *J Adv Pharm Technol Res*. 2011;2(2):81–7.
49. Kligman AM. Topical formulations in dermatology. *Arch Dermatol*. 1974; 110:213–7.
50. Brown MB, Martin GP, Jones SA. Dermal drug delivery. *Drug Deliv*. 2006;13(3):175–87.

HOW TO CITE: Ankita soni*, Atul Bisen, Rajesh Mujariya, Manjeet Singh, Formulation and Evaluation of a Herbal Analgesic Cream Containing Menthol, Camphor, and Clove Oil, Int. J. of Pharm. Sci., 2026, Vol 4, Issue 5, 2491-2499. https://doi.org/10.5281/zenodo.20126486