



**INTERNATIONAL JOURNAL OF  
PHARMACEUTICAL SCIENCES**

[ISSN: 0975-4725; CODEN(USA):IJPS00]

Journal Homepage: <https://www.ijpsjournal.com>



**Research Article**

**Phytochemical Screening And GC-MS Analysis Of *Ocimum Sanctum*,  
*Amaranthus Viridis*, And Cystone**

**Vishal Sambhaji Sutar<sup>1\*</sup>, Nitin Anandrao Kamble<sup>2</sup>**

<sup>1</sup>Research Scholar, Department of Zoology, Shivaji University, Kolhapur, Maharashtra, India. 416 004.

<sup>2</sup>Professor, Department of Zoology, Shivaji University, Kolhapur, Maharashtra, India. 416 004.

**ARTICLE INFO**

Received: 22 Aug 2024

Accepted: 26 Aug 2024

Published: 05 Sep 2024

**Keywords:**

Phytochemical screening,  
*Ocimum sanctum*,  
*Amaranthus viridis*, Cystone,  
GC-MS.

**DOI:**

10.5281/zenodo.13737965

**ABSTRACT**

In the field of pharmacology, medicinal plants have of great importance because of their therapeutic application which are natural, biocompatible, without any side effects with cost effective. The present study was carried out to study Phytochemical screening and GC-MS analysis of ethanolic extract of seeds of *Ocimum sanctum* (OS), *Amaranthus viridis* (AV) and Cystone (CS). *Ocimum sanctum* and *Amaranthus viridis* belongs to family Labiatae and Amaranthaceae respectively. Whereas Cystone is ayurvedic herbal formula made by Himalaya healthcare. Soxhlet extraction of grinded seeds was done by using absolute ethanol as a solvent. The extract was subjected to various photochemical protocols and GCMS analysis. The phytochemical screening of ethanolic extract of Seeds of *Ocimum sanctum*, *Amaranthus viridis* and Cystone revealed the presence of flavonoids, tannin, phenols, saponins, alkaloids, proteins, carbohydrate, terpenoids, and steroids. The GC-MS analysis revealed the presence of twenty-three, fifteen, and five bioactive compounds in OS, AV and CS respectively. The phytochemical and GC-MS profiling of ethanolic extract of OS, AV and CS revealed the presence of bioactive compounds with important medicinal properties.

**INTRODUCTION**

Worldwide under health sciences and for therapeutic protocol, ancient and historical literature reported huge knowledge of diseases and medication related to information provided ayurveda and folk theories [1]. According to Botanical Survey of India, near about 46,000 species of plant were identified and reported ,

where more than 7,000 plants species were documented for variety of medicinal features [2]. Among these plants, *Ocimum sanctum* (OS) commonly known as Tulsi found as sacred plant according to Hindu religion all over India . OS most respectful herb in all over the world [3].

**\*Corresponding Author:** Vishal Sambhaji Sutar

**Address:** Research Scholar, Department of Zoology, Shivaji University, Kolhapur, Maharashtra, India. 416 004. Indore

**Email** ✉: [sutarvishal1415@gmail.com](mailto:sutarvishal1415@gmail.com),

**Relevant conflicts of interest/financial disclosures:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.



In the phyto remedial procedure about all the part like leaves, stem, flower, root, seeds or even whole plant of OS prominently used for various treatments like bronchitis, malaria, diarrhoea, dysentery, skin disease, arthritis, eye diseases, insect bites, anti-fertility, anticancer, antifungal, antimicrobial, cardioprotective, analgesic, antispasmodic and adaptogenic actions. Biochemically alcoholic extract of OS reported for numerous pharmacological activities like hypoglycaemic, immunomodulatory, antistress, analgesic, antipyretic, anti-inflammatory, anti-ulcerogenic, antihypertensive, CNS depressant, radioprotective, antitumour and antibacterial activity [4, 5]. Genus *Ocimum* belonging to family Labiatae with variety of species with different therapeutic uses including *Ocimum sanctum* L. (Tulsi), *Ocimum gratissium* (Ram Tulsi), *Ocimum canum* (Dulal Tulsi), *Ocimum basilicum* (Ban Tulsi), *Ocimum kilimandscharicum*, *Ocimum ammericanum*, *Ocimum camphora*, *Ocimum minimum* L., *Ocimum tenuiflorum* L. and *Ocimum micranthum* etc.[6]. Since ancient period, among phyto remedial herb, genus *Amaranth* has notably reported for to treat conditions like diuretic, useful in cold and cough, urinary and throat troubles and gastric problems. Seed of *amaranth* were used in hypertension, cardiovascular disease, reducing blood pressure, lowering cholesterol, also it is used in piles, blood purify and antiscorbutic [7]. Specially *Amaranthus viridis* (AV) among the group has anti-inflammatory feature used in vermifuse, diuretics and also for the treatment of kidney stones as antiurolithiatic agent [8]. In traditional Indian

medicines, Cystone is a polyherbal ayurvedic formula known for treatment of, burning micturition , urinary tract complications in pregnancy and other various renal disorders [9]. Each tablet form contains approximately 130 mg *Didymocarpus pedicellate*, 98 mg *Saxifraga ligulate*, 32 mg *Rubia cordifolia*, *Cyperus scariosus* , 32 mg *Achyranthes aspera*, 32 mg *Onosma bracteatum* , 32 mg *Vernonia cinerea*, 26 mg Purified Shilajeet and 32 mg Hajrul yahood Bhasma [10]. By taking account of available phytomedicinal literature and pathological scenario present investigation was focused to find out comparative efficiency and biochemical screening of main bioactive compounds from two different plant species, pertaining to understand and upgrade traditional knowledge for its better application against some urolithiatic condition.


## **MATERIALS AND METHODS :**

### **Plant Sample:**

The best quality and non infected seeds of *Ocimum sanctum* and *Amaranthus viridis* was collected from the local market from Kolhapur, Maharashtra, India. The tablet Cystone (Himalaya, Batch No.106221154 ) was procured from local pharmacy.

### **Crude Extraction :**

Among the collected seeds 300 gm of the seeds of *Ocimum sanctum* (OS) and *Amaranthus viridis* (AV) were cleaned and grinded in mortar and pestle followed by Cystone. Absolute ethanol was used as solvent for Soxhlet extraction method for 6 hr. The prepared seed extract was evaporated and dry under reduced pressure at 40° C [11].



**Figure 1 :** Fig . 1A Whole plant of *Oscimum sanctum* (OS) ( Tulsi ) , Fig. 1B - Freshly collected noninfected, normal, well developed inflorescence of OS. , Fig. 1C - Dry healthy seeds Of OS. Fig. 1D and 1E - Morphometric feature of mature seeds of OS. (50 X) Fig. 1F and 1G - Grinded powder form of seeds OS. Fig. 1H - Assembly Soxhlet apparatus , Fig. 1I - Thimble , Fig . 1J -Ethanolic extract of seeds of *Oscimum sanctum* , Fig . 1K - GC MS Assembly used for analysis.

### GCMS Analysis protocol:

The ethanolic extracts obtained from seeds of *Ocimum sanctum*, *Amaranthus viridis* and *Cystone* were subjected to GC-MS analysis. GC-MS analysis of this extract was performed using a Shimadzu GP 2010 system and Gas Chromatograph interfaced to a Mass Spectrometer (GC-MS) equipped with a RTX TQ 2010 column (60 m X 0.25 mm ID X 1 iMdf, composed of 100% D imethyl polysiloxane) for GC-MS detection, an electron ionization system with ionization energy of 70 eV was used. Specially pure Helium gas

(99.99 %) was used as the carrier gas at a constant flow rate of 1ml/min and an injection volume of 2  $\mu$ l was employed. Injector temperature 250°C; Ion-source temperature 280° C. The oven temperature was programmed from 80°C (isothermal for 2 min.) with an increase of 10°C/min to 200° C, then 5°C/min to 280°C, ending with a 9 min. isothermal at 2800° C. Mass spectra were taken at 70 eV; a scan interval of 0.5 seconds and fragments from 45 to 650 m/z. Total GC running time was 34 minutes the relative percentage amount of each component was

calculated by comparing its average peak area to the total areas. Software adapted to handle mass spectra and chromatograms was GC-MS Real Time Analysis [12].

#### **Identification of Phytochemical compounds :**

Phytochemical screening for active compounds was performed using standard biochemical protocols as -

##### **1. Alkaloids by Dragendorff's test:**

Detection of alkaloids was done by using Dragendorff's reagent. The reagent was prepared by adding 0.85 gram of basic Bismuth nitrate in 10 ml acetic acid followed by adding 40 ml distilled water and named as solution A. 8 gm of Potassium iodide was added in 20 ml distilled water and named as B. 5ml from solution A and B was added in 20 ml of Acetic acid followed by 100 ml of water and named as C or Dragendorff's reagent [13]. 4 to 5 drops of Dragendorff's reagent was added in the 1ml of extract in a test tube. A reddish-brown precipitate was observed which indicated the presence of alkaloids [14].

##### **2. Flavonoids by Shinoda test:**

1 ml of extract was added with Magnesium followed by few drops of concentrated Hydrochloric acid. Appearance of reddish to pink colour indicates the presence of flavonoids [15].

##### **3. Glycosides by Liebermann's test.:**

2 ml of acetic acid and 2 ml of Chloroform was added in 2 ml of extract followed by concentrated Sulphuric acid. Appearance of green colour showed presence of glycosides [16]

##### **4. Proteins by Biuret test:**

Few drops of extract was added in the 1 ml 3% Copper sulphate followed by few drops of 10 % Sodium hydroxide. Appearance of violet or red colour formation indicating that proteins are present [17].

##### **5. Saponin by Foam test :**

few drops extract was added in the few drops of water . Foam produced on shaking and persists for 10 to 15 min, indicates presence of Saponins [18].

##### **6. Phytosterols by Sulphuric acid test :**

2 ml of extract was added in 2 ml Chloroform followed by concentrated Sulphuric acid. The solution was diluted in acetic acid. Finally 3 ml Acetic anhydride was added. Appearance of bluish green color showed the presence of phytosterols [19]

##### **7. Tannin by Ferric chloride test :**

2 ml of extract was diluted in distilled water. Few drops of Ferric chloride was added . Blue-green or black coloration indicates presence of tannins [20]

##### **8. Carbohydrates by Benedict's Test :**

Detection of carbohydrate was done by using Benedict's reagent. The reagent was prepared by adding 173 gm of Sodium citrate in 100 gm of Sodium carbonate and diluted with 800 ml distilled water and boiled and named as solution A. 17.3 gm of Copper sulphate dissolved in 100 ml distilled solution B. 10 ml from solution A and B was named as C or Benedict's reagent [21]. 2 ml of extract was added in 2 ml of Benedict's reagent and boiled. Appearance of reddish brown precipitate indicating the presence of carbohydrate (Reducing sugar) [22].

##### **9. Phenol by Ferric chloride test :**

1ml of the extract was added with three drops of Ferric chloride followed by few drops of Potassium ferricyanide. Appearance of greenish-blue confirmed the presence of phenols [ 23].

##### **10. Terpenoids by Salkowski test :**

2 ml of extract was mixed with 2 ml Chloroform followed by 3 ml concentrated H<sub>2</sub>SO<sub>4</sub>. appearance of reddish brown colour of indicating presence of terpenoids [24].

#### **RESULTS:**

Phytochemical screening of seed extract of *Ocimum sanctum* showed presence of flavonoids, proteins, saponin, phytosterols, carbohydrates, phenol and terpenoids. The alkaloid, glycosides and tannins were found absent. The Seed extract of *Amaranthus viridis* showed presence of alkaloid, flavonoids, proteins, saponins, tannins,




carbohydrates and phenols where as glycosides, phytosterols and terpenoids were absent. Cystone extract showed presence of alkaloids, proteins, saponins, carbohydrates and phenols where flavonoids, glycosides, phytosterols, tannins and tannin were found absent. The GC – MS chromatogram of ethanolic extract of *Ocimum sanctum*, *Amaranthus viridis* and Cystone showed many major peaks ( graph 2 a. 3 a and 4 a). The chromatogram of *Ocimum sanctum* extract found were total 23 chemical constituents as 2-Hepten-4-ol, 2-Decenal, (E) -, 2,4-Decadienal, (E,E)-, 1-Tetradecanol, Tetradecane, Caryophyllene, Benzene, 1-(1,5-dimethyl-4-hexenyl)-4-methyl, (1S,5S)-2-Methyl-5-(1,5R-6-methylhept-5en-2), 1H-Benzocycloheptene,2, 4a, 5, 6,7, 8, 9, 9a -octa, Cyclohexene, 3-(1, 5-dimethyl-4-hexenyl)-6m, 1-Hexadecanol, Hexadecane, 1,1':4'1''-Tercyclohexane, Khusimyl methyl ether, Tetradecanoic acid, 1-Nonadecene, Heneicosane, Eicosanal, i-Propyl 12- methyltetradecanote, n-

Hexadecanoic acid, Linoleic acid ethyl ether, 9,12,15- Octadecatrienoic acid and ethyl ester. Out of 23 , Hexadecenoic acid (23 %), i-Propyl 12- methyltetradecanote ( 14 %) found maximum. The chromatogram of *Amaranthus viridis* extract found total 15 chemical constituents as 1-Tetradecanol, Tetradecane, 1-Nonadecene, Hexadecane, Cyclohexane decyle, Eicosane, Dodecylcyclohexane, n-Hexadecanoic acid,, Hexadecanoic acid, ethyl ester, oleic acid, linoleic acid ethyle ester, ( E )-9-Octadecenoic acid ,15-methyl-, ethyl ester and Ethyl 14- methyl-hexadecanoate in which Hexadecenoic acid ( 21%) and linoleic acid ethyle ester ( 20 %) found maximum The chromatogram of Cystone extract found total 5 chemical constituents as (3ar,4R,7R)-1, 4, 9, 9 -1 tetramethyl-3, 4, 5, 6, 7, 8, n-Hexadecanpic acid,6- Ocadecenoic acid, Octadecanoic acid and Octadecanoic acid,17 methyl-,methyl ester where hexadecenoic acid ( 54 %) showed maximum area

**Table 1. Preliminary phytochemical analysis of ethanolic extract of seeds of *Oscimum sanctum* , *Amaranthus viridis* and Tablet Cystone.**

Sr. No.	Phytochemical constituents	Test	Result		
			O. sanctum	A. viridis	Cystone.
1	Alkaloid	Dragendorff's test	-	+	+
2	Flavonoids	Shinoda test	+	+	-
3	Glycoside	Liebermann's test	-	-	-
4	Proteins	Biuret test	+	+	+
5	Saponins	Foam test	+	+	+
6	Phytosterols	Sulphuric acid test	+	-	-
7	Tannins	Ferric chloride test	-	+	-
8	Carbohydrates	Benedict's test	+	+	+
9	Phenols	Ferric chloride test	+	+	+
10	Terpenoids	Salkowski test	+	-	-


+ : Present , - :Absent.



**Figure 2 . Graphical representation of phytochemical analysis of ethanolic extract of seeds of *Oscimum sanctum* , *Amaranthus viridis* and Cystone**

3.2 - GC-MS analysis of ethanolic extract of seeds of *Oscimum sanctum* , *Amaranthus viridis* and Cystone .

3.2.1 - GC-MS analysis of *Oscimum sanctum* :



**Figure 3: GC-MS spectrum of *Oscimum sanctum***




Figure 4: Graphical representation of percentage of area detected by GC -MS in seeds of Oscimum sanctum.

Table No. 2 : Component identified in the seed extract of Oscimum sanctum.

Sr. No.	Retention Time (min)	Name of the component	Molecular formula	Molecular weight gm/mol	Peak Area %
A	9.580	2-Hepten-4-ol	C7H14O	114	0.43
B	9.874	2-Decenal,(E)-	C10H18O	154	1.06
C	10.406	2,4-decadienal,(E,E)-	C10H16O	152	4.43
D	11.674	1-Tetradecanol	C15H30O	214	1.51
E	11.780	Tetradecane	C14H30	198	2.25
F	12.025	Caryophyllene	C14H30	198	2.89
G	12.092	Caryophyllene	C15H24	204	1.76
H	12.921	Benzene, 1-(1,5-Dimethyl)-4-methyl	C15H22	202	2.29
I	13.103	(1S,5S)-2-Methy-5(R)-6-Methyhept-5-en-2	C15H24	204	0.76
J	13.287	1H-Benzocycloheptene,2,4a,5,6,7,8,9,9a-octa	C15H24	204	1.03
K	13.527	Cyclohexene,3-(1,5-Dimethyl-4-hexenyl)-6-m	C15H24	204	1.70
L	14.353	1-Hexadecanol	C16H34O	242	5.63
M	14.442	Hexadecane	C16H34	226	6.79
N	15.252	1,1'.4',1''-Tercyclohexane	C18H38	248	5.42
O	15.909	Khusimyl methyl ether	C16H26O	234	1.41
P	16.802	Tetradecanoic acid	C14H28O2	228	4.74
Q	17.098	1-Nonadecene	C19H38	266	5.09
R	17.200	Heneicosane	C21H44	296	2.98
S	17.580	Eicosanal	C20H40O	296	1.43
T	17.992	i-Propyl 12- Methyltetradecanoate	C14H36O2	284	13.98
U	20.873	n-Hexadecanoic acid	C16H32O2	256	23.44
V	27.084	Linoleic acid Ethyl ester	C20H36O2	308	4.30
W	27.246	9,12,15-Octadecatrienoic acid, ethyl ester	C20H34O2	306	4.68

### 3.2.2 - GC-MS analysis of *Amaranthus viridis*




Figure 5: GC-MS spectrum of *Amaranthus viridis*.




Figure 6 : Graphical representation of percentage of area detected by GC -MS in seeds of *Amaranthus viridis*.

Table No. 3 : Component identified in the seed extract of *Amaranthus viridis*.

Sr. No.	Retention Time (min)	Neme of the component	Molecular formula	Molecular weight gm/mol	Peak Area %
A	11.681	1-Tetradecanol	C <sub>14</sub> H <sub>30</sub> O	214	0.47
B	11.786	Tetradecane	C <sub>14</sub> H <sub>30</sub>	198	0.81
C	14.357	1-Nonadecane	C <sub>19</sub> H <sub>38</sub>	266	3.62
D	14.445	Hexadecane	C <sub>16</sub> H <sub>34</sub>	226	3.09
E	15.226	Cyclohexane	C <sub>6</sub> H <sub>12</sub>	84	1.39
F	17.102	1-Nonadecane	C <sub>19</sub> H <sub>38</sub>	266	3.80
G	17.207	Eicosane	C <sub>20</sub> H <sub>42</sub>	282	2.33
H	18.546	Dodecylcyclohexane	C <sub>18</sub> H <sub>36</sub>	252	0.52
I	20.909	n-Hexadecanoic acid	C <sub>16</sub> H <sub>32</sub> O <sub>2</sub>	256	14.48
J	21.480	Hexadecanoic acid, ethyl ester	C <sub>18</sub> H <sub>36</sub> O	284	20.91
K	26.853	Oleic acid	C <sub>18</sub> H <sub>34</sub> O <sub>2</sub>	282	4.13
L	27.091	Linoleic acid ethyl ester	C <sub>20</sub> H <sub>36</sub> O <sub>2</sub>	308	20.34

M	27.259	(E)-9-Octadecanoic acid ethyl ester	C20H38O2	310	17.34
N	27.917	Heptadecanoic acid, 15-methyl-, ethyle ester	C20H40O2	298	4.86
O	31.659	Ethyl 14-methyl-hesadecanoate	C18H36O2	298	1.89
A	11.681	1-Tetradecanol	C14H30O	214	0.47
B	11.786	Tetradecane	C14H30	198	0.81
C	14.357	1-Nonadecane	C19H38	266	3.62
D	14.445	Hexadecane	C16H34	226	3.09
E	15.226	Cyclohexane	C16H32	224	1.39
F	17.102	1-Nonadecane	C19H38	266	3.80
G	17.207	Eicosane	C20H42	282	2.33
H	18.546	Dodecylcyclohexane	C18H36	252	0.52

### 3.2.2 - GC-MS analysis of Cystone:




Figure 7: GC-MS spectrum of *Amaranthus viridis*




Figure 8 : Graphical representation of percentage of area detected by GC -MS in Cystone.

### DISCUSSION :

By the therapeutic point of view generally phytochemicals were reported as biologically active compounds, which protect animals from natural or induced toxic chemicals, pathogen where disease causing microbes are in association of fruits, leaves, seeds, grains herbs and some time

total plant spices [25]. Along with food nutrients, foods source, plants have rich source of bioactive phytochemicals or bio nutrients which includes preventing chronic diseases like cancer, diabetes, coronary heart disease and hypercholesterolaemia. Some group of phytochemicals have disease-preventing antioxidants,

detoxification , immunological and neuropharmacological activators [26]. In modern pharmacological and nutraceutical industries, phytochemicals played an important role because of wide range of its application like cofactors, modulators, inhibitors, antioxidant. The contents like carotenoids, catechins, curcumin, diosgenin , polyphenol and flavonoids have wide range against treatment of various human disease [27]. [28] documented various beneficiary effects of flavonoid like antioxidant, anti-inflammatory, antiallergic, anti-microbial, effective in hepatotoxicity, cardiovascular diseases, gastric ulcer, rheumatic disease, thrombosis, memory cognition and in cardiovascular disease. In the present study ethanolic extract of seed of *Ocimum sanctum*, *Amaranthus viridis* and Cystone, the GC MS analysis conforms variety of pharmacogenic constituents like, phytol, octadecanoic acid, Palmitic acid, oleic acid common in all three seed extract. Also Cetane and isocaryophyllene were identified. Along with precursor for vitamin E and vitamin K, phytol showed anxiolytic, metabolism-modulating, cytotoxic, antioxidant, autophagy and apoptosis inducing, antinociceptive, anti-inflammatory, immune-modulating, and antimicrobial effects [29, 30]. [31] reported 9, 12-Octadecadienoic acid has the property of antioxidant, anti-inflammatory and antiarthritic in his work. Cetane number is a indicators of the quality of diesel fuel. Variety of feedstock vegetables oils with K45 – K67 are useful for the production of biodiesel [32]. [ 33] reported  $\beta$ -caryophyllene has antioxidant, anti microbial and anti bacterial activities. [ 34.] reported phenols and alkaloids have antioxidants properties. [35] reported flavonoid, alkaloid, phenol, tannin and saponin have significant level of antioxidant.

#### CONCLUSION :

On the basis of result obtained under present investigation study revealed that, comparatively

extract of seeds of *Ocimum sanctum* , *Amaranthus viridis* and Cystone has variety of bioactive compound which were screened by assay including flavonoids, alkaloids, phenols , tannin, saponin, phytol, etc. All these compound known for biological role in the animal body as antioxidant, antimicrobial, anticancer, anti-ulcer, anti-inflammatory and hepatoprotective. Comparatively OS, AV and CS has hexadecenoic acid chemical compound found prominently which play important role in in dissociation of calcium carbonate content. Both the phytoextract were seems to be useful for the protection of cells from accumulation of crystals and relevant pathological condition in comparison to cystone. The further advance study such as qualitative determination, purification and characterization of phytochemical constituents all extract may help to formulate herbal preparation of medical use.

#### ABBREVIATIONS

GC-MS : Gas Chromatography Mass Spectrometry.

OS : *Ocimum sanctum*

AV : *Amaranthus viridis*

CS : Cystone

CNS : Central Nervous System

mg : Milli Gram

ml : Milli litter

gm : Gram

CFC : Central Facility Centre

°C : Degree Celsius

eV : Electron ionization

m/z : Mass to Charge ratio

DL : Detection limit.

H<sub>2</sub>SO<sub>4</sub> : Sulphuric acid

+ : Present

- : Absent

gm/mol : Gram per Molecule

#### ACKNOWLEDGEMENT

Authors are thankful to Head, Department of Zoology, Shivaji University, Kolhapur for support during the present research work.



## CONFLICT OF INTEREST

There are no conflicts of interest.

## REFERENCES

1. Vaidya AD, Devasagayam TP (2007) Current status of herbal drugs in India: an overview. *J Clin Biochem Nutr.* Jul;41(1):1-11. doi: 10.3164/jcbrn.2007001. PMID: 18392106; PMCID: PMC2274994.
2. Bisoi SS & Panda D (2015). Ethno-Medicinal Plants Present In Sacred Groves Of Koraput District Of Odisha, India. *Acta Biomedica Scintia.* 2. 39-42.
3. Kulkarni KV, Belvotagi VA (2018) A review on: Indian traditional shrub Tulsi (*Ocimum sanctum*): The unique medicinal plant, *Journal of Medicinal Plants Studies* 2018; 6(2): 106-110.
4. Pattanayak P, Behera P, Das D, Panda SK (2010 ) *Ocimum sanctum* Linn. A reservoir plant for therapeutic applications: An overview. *Pharmacogn Rev.* 2010 Jan;4(7):95-105. doi: 10.4103/0973-7847.65323. PMID: 22228948; PMCID: PMC3249909.
5. Das SK & Vasudevan D (2006) Tulsi: The Indian Holy power Plant. *Natural Product Radiance.* 5. 279-283.
6. Das AK , Hossain M & Deb N (2010). Chemical Compositions of Different Extracts of *Ocimum basilicum* Leaves. *Journal of Scientific Research.* 3. 197-206. 10.3329/jsr.v3i1.5409.
7. Rastogi A, Shukla S. (2013) Amaranth: a new millennium crop of nutraceutical values. *Crit Rev Food Sci Nutr.* 2013;53(2):109-25. doi: 10.1080/10408398.2010.517876. PMID: 23072528.
8. Asha S, & Thirunavukkarasu P (2013). Antiuro lithiatic Activity of *Amaranthus viridis* on Ethylene Glycol Induced Male Rats. 2013. 13-17.
9. Rao M, Rao MN ( 1998) Protective effects of cystone, a polyherbal ayurvedic preparation, on cisplatin-induced renal toxicity in rats, *Journal of Ethnopharmacology*, Volume 62, Issue 1, 1998,Pages 1-6, ISSN 0378-8741, [https://doi.org/10.1016/S0378-8741\(98\)00003-8](https://doi.org/10.1016/S0378-8741(98)00003-8).
10. Mohanty N.K. & Nayak, R.L. & Patki, Pralhad. (2010). Safety and Efficacy of an Ayurvedic Formulation Cystone in Management of Ureteric Calculi: A Prospective Randomized Placebo Controlled Study. *American Journal of Pharmacology and Toxicology.* 5. 58-64. 10.3844/ajptsp.2010.58.64.
11. Lim SN, Cheung PC, Ooi VE, Ang PO (2002) Evaluation of antioxidative activity of extracts from a brown seaweed, *Sargassum siliquastrum*. *J Agric Food Chem.* 2002 Jun 19;50(13):3862-6. doi: 10.1021/jf020096b. PMID: 12059172.
12. Phukan H, Bora C & Mitra P. (2017) Phytochemical Screening and GC-MS Analysis of Methanolic leaf Extract of an Endemic Plant *Kayea assamica*. *IOSR Journal of Pharmacy and Biological Sciences.* 12. 7-16. 10.9790/3008-1205020716.
13. Sanghani Y (2017). Handbook of chemistry lab reagent.
14. Samatha, T. & Srinivas, P. & Shyamsundarachary, R. & Nanna RS (2012). Phytochemical analysis of seeds, stem bark and root of an endangered medicinal forest tree *Oroxylum Indicum(L)kurz*. *International Journal of Pharma and Bio Sciences.* 3. B1063-B1075.
15. Guediri, I.; Boubekri, C.; Smara, O (2020) Preliminary Phytochemical Screening From Different Extracts Of *Solanum Nigrum* Plant Growing In South Of Algeria. *Journal Of Fundamental And Applied Sciences*, [S. L.], V. 12, N. 2, P. 624–633, 2020. Doi: 10.4314/Jfas.V12i2.7.
16. Gul R, Jan SU, Faridullah S, Sherani S, Jahan N (2017) Preliminary Phytochemical



- Screening, Quantitative Analysis of Alkaloids, and Antioxidant Activity of Crude Plant Extracts from *Ephedra intermedia* Indigenous to Balochistan. *ScientificWorldJournal*. 2017;2017:5873648. doi: 10.1155/2017/5873648. Epub 2017 Mar 13. PMID: 28386582; PMCID: PMC5366796.
17. Kancherla N, Dhakshinamoothi A, Chitra K, Komaram RB (2019) Preliminary Analysis of Phytoconstituents and Evaluation of Anthelmintic Property of *Cayratia auriculata* (In Vitro). *Maedica (Bucur)*. 2019 Dec;14(4):350-356. doi: 10.26574/maedica.2019.14.4.350. PMID: 32153665; PMCID: PMC7035446.
18. Bhandary S, Kumari S & Bhat V, Sherly S & Bekal M (2012). Preliminary phytochemical screening of various extracts of *Punica granatum* peel, whole fruit and seeds. *Nitte University Journal of Health Science*. 2. 10.1055/s-0040-1703609.
19. Ahmed Z, Aziz S, Hanif M, Mohiuddin SG, Ali Khan SH, Ahmed R, Sheikh Ghadzi SM, Naoras Bitar A (2020) Phytochemical screening and enzymatic and antioxidant activities of *Erythrina suberosa* (Roxb) bark. *J Pharm Bioallied Sci*. Apr-Jun;12(2):192-200. doi: 10.4103/jpbs.JPBS\_222\_19. Epub 2020 Apr 10. PMID: 32742119; PMCID: PMC7373117.
20. Ukoha P, Egbuonu, Anthony C, Nnamdi O Ejikeme P (2011). Tannins and other phytochemical of the *Samanea saman* pods and their antimicrobial activities. *Afri. J. Pure Appl. Chem.*. 5. 237-244.
21. Shaikh J, & Patil M. (2020). Qualitative tests for preliminary phytochemical screening: An overview. 8. 603-608. 10.22271/chemi.2020.v8.i2i.8834.
22. Deshmukh M Theng M (2018). Phytochemical Screening, Quantitative Analysis Of Primary And Secondary Metabolites Of *Acacia Arabica* Bark. *International Journal of Current Pharmaceutical Research*. 10. 35. 10.22159/ijcpr.2018v10i2.25889.
23. Dubale S, Kebebe D, Zeynudin A, Abdissa N, Suleman S (2023) Phytochemical Screening and Antimicrobial Activity Evaluation of Selected Medicinal Plants in Ethiopia. *J Exp Pharmacol*. Feb 8;15:51-62. doi: 10.2147/JEP.S379805. PMID: 36789235; PMCID: PMC9922502.
24. Sheel, D., & Nisha, K.M. (2014). Qualitative phytochemical analysis for isolation of terpenes from *Clerodendron infortunatum* leaves. *IOSR Journal of Applied Chemistry*, 7, 14-18.
25. Saxena M & Jyoti S & Nema R & Dharmendra S & Abhishek, G. (2013). Phytochemistry of Medicinal Plants. *J Pharm Phytochem*. 1. 168-182.
26. Rao BN (2023) Bioactive phytochemicals in Indian foods and their potential in health promotion and disease prevention. *Asia Pac J Clin Nutr*. 2003;12(1):9-22. PMID: 12737006.
27. Singh MK, Singh SK, Singh AV, Verma H, Singh PP, Kumar A, (2020) *Phytochemicals: Intellectual property rights*, Editor(s): Bhanu Prakash, *Functional and Preservative Properties of Phytochemicals*, Academic Press, 2020, Pages 363-375, ISBN 9780128185933, <https://doi.org/10.1016/B978-0-12-818593-3.00012-9>.
28. Sandhar HK, Kumar B , Prasher S. & Tiwari P & Salhan, M. & Sharma P.(2011). A review of phytochemistry and pharmacology of flavonoids. *Internationale Pharmaceutica Scientia*. 1. 25-41.
29. Casuga F, Castillo & Corpuz MJA (2016) GC-MS analysis of bioactive compounds present in different extracts of an endemic plant *Broussonetia luzonica* (Blanco) (Moraceae) leaves. *Asian Pacific Journal of Tropical Biomedicine*. 6. 10.1016/j.apjtb.2016.08.015.

30. Islam MT , Eunüs SA , Shaikh UJ , Subrata S , Islam MA, Ahmed MI , Manik C , Utpal KK, Nagendra SY, Khan IN, Billah MM , Magdalena DP, Gokhan Z, Clemens M, Ferdinando N, Diana G, Neagoe BI , Apostol A, Maciej B, Andy Y, Demerdash A, Jianbo X, Dey P, Yele S, Jóźwik A, Strzałkowska N, Marchewka J, Rengasamy KRR, Horbańczuk J, Kamal MA, Mubarak MS , Mishra SK , Shilpi JA, Atanasov AG (2018)Phytol: A review of biomedical activities, Food and Chemical Toxicology, Volume 121, 2018, Pages 82-94, ISSN 0278-6915, <https://doi.org/10.1016/j.fct.2018.08.032>.
31. Nishanthini A, Mohan VR, Jeeva S (2014) Phytochemical, FT-IR, and GC-MS analysis of stem and leaf of *Tiliacora acuminata* (lan.) hook f and Thomas(menispermaceae). Int J Pharm Sci Res 5(9):3977–3986.
32. Yadav A & Mishra SP , Kendurkar P, Kumar A & Maurya R (2020). Physicochemical characterization of *Jatropha* oil seed and suitability as biodiesel feedstock. Tropical Plant Research. 7. 581-586. [10.22271/tpr.2020.v7.i3.071](https://doi.org/10.22271/tpr.2020.v7.i3.071).
33. Sitarek, Rijo, Patricia, Garcia, Catarina, Skala, Ewa, Kalemba, Danuta, Białas, AdamJ., Szemraj, Janusz, Pytel, Dariusz, Toma, Monika, Wysokińska, Halina, Śliwiński, Tomasz (2017)Antibacterial, Anti-Inflammatory, Antioxidant, and Antiproliferative Properties of Essential Oils from Hairy and Normal Roots of *Leonurus sibiricus* L. and Their Chemical Composition, Oxidative Medicine and Cellular Longevity, 2017, 7384061, 12 pages, 2017. <https://doi.org/10.1155/2017/7384061>
34. Gan J, Feng Y, He Z, Li X , Zhang H, ( 2017) Correlations between Antioxidant Activity and Alkaloids and Phenols of Maca (*Lepidium meyenii*), Journal of Food Quality, 2017, 3185945, 10 pages, 2017. <https://doi.org/10.1155/2017/3185945>
35. Nithya TG , Jayaprakash J, Raguathan M (2016). Antioxidant activity, total phenol, flavonoid, alkaloid, tannin, and saponin contents of leaf extracts of *Salvinia molesta* D. S. Mitchell (1972). 9.

**HOW TO CITE:** Vishal Sambhaji Sutar, Nitin Anandrao Kamble , Phytochemical Screening And GC-MS Analysis Of *Ocimum Sanctum*, *Amaranthus Viridis*, And Cystone, Int. J. of Pharm. Sci., 2024, Vol 2, Issue 9, 258-270. <https://doi.org/10.5281/zenodo.13737965>

